

Efficacy of systemic enzyme therapy as an anti-inflammatory agent when used as monotherapy or in combination with NSAIDs: A systematic review

Ganiga Channaiah Shivakumar^{1,A,F}, Nishath Sayed Abdul^{2,B,C}, Mahesh Shenoy^{2,C,D}, Abdullah Abdulaziz Saeidi^{3,C}, Vincenzo Ronsivale^{4,C,D}, Hande Uzunçibuk^{5,D,E}, Gabriele Cervino^{4,E,F}, Maria Maddalena Marrapodi^{6,7,E,F}, Giuseppe Minervini^{8,E,F}

¹ Department of Oral Medicine and Radiology, People's College of Dental Sciences and Research Centre, People's University, Bhopal, India

² Department of Oral Maxillofacial Surgery and Diagnostic Sciences, Faculty of Oral Pathology, College of Medicine and Dentistry, Riyadh Elm University, Saudi Arabia

³ Department of Diagnostic Oral and Maxillofacial Sciences, College of Dentistry, Taibah University, Medina, Saudi Arabia

⁴ Department of Biomedical and Dental Sciences, Morphological and Functional Images, University of Messina, G. Martino Polyclinic, Messina, Italy

⁵ Department of Orthodontics, Faculty of Dentistry, Trakya University, Edirne, Turkey

⁶ Saveetha Dental College and Hospitals, Saveetha Institute of Medical and Technical Sciences (SIMATS), Chennai, India

⁷ Department of Woman, Child and General and Specialist Surgery, University of Campania Luigi Vanvitelli, Naples, Italy

⁸ Multidisciplinary Department of Medical-Surgical and Dental Specialties, University of Campania Luigi Vanvitelli, Naples, Italy

A – research concept and design; B – collection and/or assembly of data; C – data analysis and interpretation;

D – writing the article; E – critical revision of the article; F – final approval of the article

Dental and Medical Problems, ISSN 1644-387X (print), ISSN 2300-9020 (online)

Dent Med Probl.

Address for correspondence

Giuseppe Minervini

E-mail: giuseppe.minervini@unicampania.it

Funding sources

None declared

Conflict of interest

None declared

Acknowledgements

The authors would like to thank Prof. Luis Eduardo Almeida, a native English speaker and associate professor in the Department of Surgical Sciences at the School of Dentistry, Marquette University (Milwaukee, USA), for reviewing and editing the manuscript to ensure appropriate scientific English.

Received on February 12, 2025

Reviewed on May 12, 2025

Accepted on June 8, 2025

Published online on October 28, 2026

Cite as

Shivakumar GC, Abdul NS, Shenoy M, et al. Efficacy of systemic enzyme therapy as an anti-inflammatory agent when used as monotherapy or in combination with NSAIDs: A systematic review [published online as ahead of print on October 28, 2026]. *Dent Med Probl.* doi:10.17219/dmp/206041

DOI

10.17219/dmp/206041

Copyright

Copyright by Author(s)

This is an article distributed under the terms of the Creative Commons Attribution 3.0 Unported (CC BY 3.0) (<https://creativecommons.org/licenses/by/3.0/>)

Abstract

Systemic enzyme therapy (SET) has emerged as a potential anti-inflammatory intervention, either as monotherapy or in combination with non-steroidal anti-inflammatory drugs (NSAIDs). Despite its increasing clinical application, the comparative efficacy of SET in various inflammatory conditions remains insufficiently explored.

A systematic review was conducted in accordance with the PRISMA (Preferred Reporting Items for Systematic reviews and Meta-Analyses) guidelines to evaluate the effectiveness of SET as an anti-inflammatory therapy. Databases were searched for studies meeting predefined eligibility criteria, including randomized and non-randomized trials. Study characteristics, intervention protocols and outcomes were extracted and analyzed. The risk of bias was assessed using the ROBINS-I (Risk Of Bias In Non-randomized Studies of Interventions) and Cochrane risk-of-bias (RoB) 2.0 tools.

Eleven studies with sample sizes ranging from 30 to 487,372 participants were included. Systemic enzyme therapy demonstrated comparable efficacy to diclofenac in terms of pain relief and functional improvement across various conditions, including joint disorders and temporomandibular disorders (TMD). Combination therapy resulted in significantly greater pain reduction ($p < 0.05$) and improved functional outcomes, as reported in several studies. In most cases, adverse effects were mild, and the safety profile of SET was more favorable than that of NSAIDs. One study reported that the concomitant use of NSAIDs and antihypertensive medications was associated with a 31% increased risk of acute kidney injury during the first 30 days.

Systemic enzyme therapy demonstrated efficacy comparable to that of NSAIDs, with a potentially more favorable safety profile in the management of inflammatory conditions. Combination therapy may provide additional benefits in terms of pain reduction and functional improvement. However, the heterogeneity of study designs and the limited number of large-scale trials warrant further research.

Keywords: systemic enzyme therapy, anti-inflammatory therapy, NSAIDs, combination therapy, pain relief

Highlights

- The study evaluates the efficacy, safety and tolerability of systemic enzyme therapy for inflammatory conditions, particularly temporomandibular disorders and osteoarthritis.
- Systemic enzyme therapy shows efficacy comparable to that of non-steroidal anti-inflammatory drugs and may improve treatment outcomes when used in combination with them.
- Systemic enzyme therapy exhibits multifaceted anti-inflammatory mechanisms, including cytokine modulation, reduced leukocyte migration and promotion of tissue repair.
- Treatment with oral enzyme combinations may offer a safer alternative or adjunct to non-steroidal anti-inflammatory drugs, with potentially fewer gastrointestinal, renal and cardiovascular risks.
- Further high-quality, large-scale randomized controlled trials are needed to confirm these findings, standardize treatment protocols and optimize clinical use.

Introduction

Systemic enzyme therapy (SET) has attracted considerable interest for its potential anti-inflammatory effects, both as monotherapy and as an adjunct to conventional non-steroidal anti-inflammatory drugs (NSAIDs).¹ Systemic enzyme preparations, consisting of combinations of proteolytic enzymes and other active agents, such as trypsin, bromelain and the flavonoid rutin, are thought to act through multiple mechanisms, including fibrinolysis, the reduction of pro-inflammatory cytokines and the promotion of tissue repair.² Unlike NSAIDs, which primarily work by inhibiting cyclooxygenase enzymes and thereby reducing prostaglandin synthesis, SET may exert its effects through multiple pathways involving inflammatory mediators, potentially avoiding some of the gastrointestinal and cardiovascular adverse effects associated with chronic use of NSAIDs.³

The inflammatory response plays a central role in various acute and chronic conditions, including musculoskeletal injuries and autoimmune disorders. Although widely regarded as a cornerstone of anti-inflammatory therapy, long-term NSAID use is often limited by adverse effects, including gastric ulceration, renal dysfunction and an increased risk of thrombotic events.^{4,5} These limitations have prompted interest in complementary or alternative therapies, particularly those that may provide anti-inflammatory benefits with a more favorable safety profile. The proposed systemic absorption of proteolytic enzymes and their potential biochemical activities make SET a possible therapeutic approach for reducing inflammation while potentially minimizing the adverse effects associated with conventional anti-inflammatory medications.⁶

Several recently published studies indicate growing interest in SET for various inflammatory conditions, such as osteoarthritis (OA), rheumatoid arthritis and postoperative inflammation.⁷ Preliminary evidence also suggests that SET may act by degrading circulating immune complexes, modulating vascular permeability, and reducing leukocyte adhesion and migration.⁸ However, these proposed benefits have not been conclusively established, as inconsistencies in study design, sample size and outcome

measures have resulted in conflicting findings regarding the efficacy of SET. Furthermore, questions related to the bioavailability of systemic enzymes and their ability to reach inflammatory sites at therapeutically relevant concentrations continue to limit their widespread adoption in clinical practice.^{9–11}

When used in combination therapy with NSAIDs, SET is thought to enhance their anti-inflammatory effect or allow for dose reductions.^{9–12} These outcomes could potentially reduce the incidence of adverse events. However, the potential additive or synergistic effects of SET in combination with NSAIDs remain poorly understood, and high-quality evidence is insufficient to draw definite conclusions. Furthermore, the lack of standardized protocols regarding enzyme formulations, dosages and treatment duration complicates direct comparisons across studies and highlights the need for strict evaluation. Given these knowledge gaps, this systematic review aims to critically evaluate the available evidence regarding the efficacy of SET as an anti-inflammatory intervention, both as monotherapy and in combination with NSAIDs.

Material and methods

PECOS framework and PRISMA compliance

The PECOS framework for this review was developed based on the PRISMA (Preferred Reporting Items for Systematic reviews and Meta-Analyses) 2020 reporting guidelines,¹³ which ensure transparent and standardized reporting. The framework was defined as follows: Population (P) – patients diagnosed with inflammatory conditions, including joint disorders and temporomandibular disorders (TMD); Exposure (E) – SET administered as monotherapy or in combination with NSAIDs; Comparison (C) – placebo or NSAID treatment; Outcome (O) – reductions in inflammation and pain, improvements in joint mobility, and patient-reported functional scores; Study design (S): randomized controlled trials (RCTs), non-RCTs and cohort studies.

Inclusion and exclusion criteria

The inclusion criteria required studies to investigate patients with inflammatory conditions treated with SET as monotherapy or in combination with NSAIDs. Eligible studies included RCTs, non-RCTs and cohort studies involving human populations. Studies were required to report at least 1 measure of inflammation reduction, pain relief or functional improvement. Only full-text articles published in English were considered, with no restrictions on the year of publication.

Studies involving in vitro or animal models, non-inflammatory conditions or SET formulations that did not include trypsin, bromelain or rutoside were excluded. Reviews, editorials and articles lacking sufficient methodological details or reporting outcomes beyond the scope of this review were also excluded. Conference abstracts were excluded unless they reported sufficient methodological and outcome data. Studies published in languages other than English were not considered.

Database search

A systematic literature search was performed across 6 databases: PubMed; Embase; the Cochrane Library; Scopus; Web of Science; and CINAHL (Cumulative Index to Nursing and Allied Health Literature). The search terms included “systemic enzyme therapy”, “anti-inflammatory”, “diclofenac”, “bromelain”, “trypsin”, “rutoside”, “joint health”, “temporomandibular joint”, and “inflammation”. Boolean operators, such as AND, OR and NOT, were used to refine search results and identify relevant studies. MeSH terms were combined with free-text terms to improve search sensitivity (Table 1).

Data extraction protocol and items

The study used a predesigned template for data extraction. The following information was extracted: study characteristics, such as author names, publication year, country,

and study design; population details, including sample size, demographics and the condition being treated; intervention details, such as formulation, dosage and treatment duration; comparator information; and outcome measures, including pain reduction, inflammatory markers, functional scores, and adverse events. Additional information included funding sources, disclosure of conflict of interest and risk-of-bias assessments.

Risk-of-bias assessment

The ROBINS-I (Risk Of Bias In Non-randomized Studies of Interventions)¹⁴ was used to assess the risk of bias in non-randomized studies, including bias due to confounding, selection of participants, classification of interventions, and measurement of outcomes. For RCTs, the Cochrane risk-of-bias (RoB) 2.0 tool¹⁵ was used to evaluate bias arising from the randomization process, deviations from the intended interventions, missing outcome data, and selective reporting. Risk-of-bias assessments were performed independently by 2 reviewers, and any disagreements were resolved through consensus.

Results

Study selection process

A total of 381 records were identified through database searches, of which 46 duplicates were removed before screening. The remaining 335 records were screened for relevance, and none were excluded at this stage. Subsequently, 37 reports could not be retrieved, leaving 298 reports for the assessment of eligibility. Of these, reports were excluded for the following reasons: failure to meet the PECOS eligibility criteria ($n = 68$); in vitro study design ($n = 48$); case reports ($n = 76$); literature reviews ($n = 55$); and theses ($n = 40$). Ultimately, 11 studies^{16–26} met the eligibility criteria and were included in the final review (Fig. 1).

Table 1. Search strategy used in the systematic review

Database	Search string
PubMed	("systemic enzyme therapy"[MeSH Terms]) AND ("anti-inflammatory agents"[MeSH Terms] OR "diclofenac"[All Fields]) AND ("joint health"[MeSH Terms] OR "temporomandibular joint disorders"[MeSH Terms])
Embase	('systemic enzyme therapy' OR 'bromelain' OR 'trypsin' OR 'rutoside') AND ('anti-inflammatory' OR 'diclofenac') AND ('joint health' OR 'temporomandibular disorders')
Cochrane Library	("systemic enzyme therapy" AND "anti-inflammatory") OR ("bromelain" AND "diclofenac") AND ("joint health" OR "temporomandibular joint")
Scopus	TITLE-ABS-KEY(("systemic enzyme therapy" OR "bromelain" OR "trypsin" OR "rutoside") AND ("anti-inflammatory" OR "diclofenac") AND ("joint health" OR "temporomandibular joint"))
Web of Science	TS=("systemic enzyme therapy" AND "bromelain" OR "trypsin" OR "rutoside") AND ("anti-inflammatory agents" OR "diclofenac") AND ("joint health" OR "temporomandibular joint disorders")
CINAHL	("systemic enzyme therapy" OR "bromelain" OR "trypsin" OR "rutoside") AND ("anti-inflammatory agents" OR "diclofenac") AND ("joint health" OR "temporomandibular joint disorders")

CINAHL – Cumulative Index to Nursing and Allied Health Literature.




Fig. 1. PRISMA (Preferred Reporting Items for Systematic reviews and Meta-Analyses) flow diagram illustrating the study selection process

Publication year and geographical distribution

The studies covered a wide time span, ranging from 1998¹⁹ to 2020 (Table 2).¹⁸ Regarding geographical distribution, a substantial proportion of studies were conducted in Germany,^{16,17,21,22,25,26} reflecting considerable research activity in this country. Other countries represented were India,^{18,20} Mexico¹⁹ and the UK.²³ The geographical diversity provided insights into the application of SET across different patient populations and healthcare settings.

Study design

Most studies employed randomized, double-blind, parallel-group designs,^{16,19,21,22,25} which are generally considered the gold standard for minimizing bias in clinical trials. A placebo-controlled trial was conducted in 2015,¹⁷ while a retrospective cohort analysis was performed in the UK in 2013.²³ Two studies employed single-blind or comparator-controlled designs,^{20,26} with slight variations in methodology tailored to the specific clinical conditions investigated.

Sample size

Sample sizes varied considerably between studies, ranging from small cohorts of 30 participants in India^{18,20} to a large retrospective cohort study involving 487,372 participants

in the UK.²³ Most of the randomized trials included between 50 and 268 participants,^{16,17,19,21,22,25,26} providing a reasonable balance between statistical power and feasibility.

Duration of interventions

Intervention durations and follow-up periods varied from 10 days²⁰ to a follow-up of 5.9 years in a retrospective study.²³ Most of the randomized trials lasted between 3 and 6 weeks,^{16,19,21,22,24,25} while one extended to 12 weeks,¹⁷ and others included additional follow-up assessments.^{25,26} These differences in duration reflected variations in the clinical conditions investigated and the study objectives.

Focus areas

The studies primarily focused on joint disorders, particularly those affecting the knee.^{16,17,19,21,24–26} One study also investigated hip-related conditions.²² Two studies targeted TMD, specifically internal derangement and OA of the temporomandibular joint (TMJ).^{18,20} Another distinct area of investigation was the risk of acute kidney injury associated with the concomitant use of NSAIDs and antihypertensive medications.²³ The diversity of conditions investigated reflects the broad range of clinical contexts in which SET has been evaluated.

Key findings

Several studies reported that SET demonstrated efficacy comparable to that of diclofenac in reducing pain and inflammation.^{16,17,20–22,25,26} Additional findings included a safety profile comparable to that of placebo¹⁷ and long-lasting therapeutic effects even after treatment.²⁵ The combination of enzymes with diclofenac was associated with improved clinical outcomes, including greater pain reduction, enhanced joint function and improved masticatory function in patients with TMD.^{18,20} A large retrospective study reported a 31% increased risk of acute kidney injury associated with triple therapy involving NSAIDs and antihypertensive agents, particularly during the first 30 days.²³ Collectively, these findings suggest that SET is effective and may reduce the risk of adverse effects, particularly when used in combination with conventional anti-inflammatory therapy.

Risk-of-bias assessment

Most RCTs demonstrated a low risk of bias across several key domains, particularly those related to the randomization process, deviations from intended interventions and outcome measurement.^{17,18,21,22,25} Additionally, some concerns regarding outcome measurement and selective reporting influenced overall ratings.^{19,24,26} Despite these reservations, the vast majority of the RCTs were assessed as having an overall low risk of bias (Fig. 2).

Table 2. Characteristics and key findings of the studies included in the review

Study	Study design	Country	Sample size, <i>n</i>	Duration	Focus area	Key findings
Akhtar et al. 2004 ¹⁶	randomized, double-blind, parallel-group comparator trial	Germany	103	6 weeks	knee disorders	OEC demonstrated efficacy comparable to that of diclofenac.
Bolten et al. 2015 ¹⁷	randomized, double-blind, placebo-controlled comparator trial	Germany	150	12 weeks	knee disorders	OEC demonstrated efficacy comparable to that of diclofenac, with reduced use of rescue medication and a tolerability profile similar to that of placebo.
Gupta et al. 2020 ¹⁸	randomized case-control clinical trial	India	30 (15 per group)	14 days	internal derangement of the TMJ	Combination therapy significantly reduced pain and joint sounds and improved chewing ability and mandibular movements. No significant between-group differences were observed in mouth opening.
Herrera 1998 ¹⁹	randomized, double-blind, parallel-group comparator trial	Mexico	59	3 weeks	monoarticular gonarthrosis (knee)	The LAFI scores and composite symptom scores were comparable between treatment groups at the end of the study. However, the sample size was insufficient to statistically establish equivalence.
Jayachandran and Khobre 2017 ²⁰	randomized, comparator-controlled trial	India	30	10 days	TMJ osteoarthritis	OEC demonstrated efficacy comparable to that of diclofenac. Combination therapy resulted in significantly greater pain reduction ($p < 0.05$) than either treatment alone.
Klein and Kullich 2000 ²¹	randomized, double-blind, parallel-group comparator trial	Germany	73	3 weeks	knee disorders	OEC demonstrated efficacy comparable to that of diclofenac.
Klein et al. 2006 ²²	randomized, double-blind, parallel-group comparator trial	Germany	90	6 weeks	hip disorders	OEC demonstrated efficacy comparable to that of diclofenac regarding the WOMAC domains of pain, stiffness and physical function, the LAFI score, investigator- and patient-reported assessments of efficacy, and responder rates based on pain and physical function.
Lapi et al. 2013 ²³	retrospective cohort study with a nested case-control analysis	United Kingdom	487,372	mean follow-up: 5.9 years	risk of acute kidney injury associated with concomitant NSAID and antihypertensive use	Triple therapy involving diuretics, ACE inhibitors/ARBs, and NSAIDs was associated with a 31% increased risk of acute kidney injury, particularly during the first 30 days of treatment.
Roth and Stauder 2001 ²⁴	randomized, double-blind, parallel-group comparator trial (conference poster)	not specified	268	6 weeks	knee disorders	Diclofenac was modestly superior on the LAFI ($MD: 1.10$; 95% $CI: 0.24-1.96$), but the difference remained within the 3-point non-inferiority margin, while 42.9% of participants were excluded from the per-protocol analysis.
Singer et al. 2001 ²⁵	randomized, double-blind, parallel-group comparator trial	Germany	63	3 weeks of treatment followed by a 4-week observation period	knee disorders	OEC demonstrated efficacy comparable to that of diclofenac, with therapeutic effects persisting after the end of treatment.
Tilwe et al. 2001 ²⁶	randomized, single-blind, parallel-group comparator trial	Germany	50	3 weeks of treatment followed by a 4-week observation period	knee disorders	OEC demonstrated overall efficacy comparable to that of diclofenac, with significantly greater improvement in joint tenderness.

ACE – angiotensin-converting enzyme; ARB – angiotensin receptor blocker; *CI* – confidence interval; LAFI – Lequesne algofunctional index; *MD* – mean difference; NSAID – non-steroidal anti-inflammatory drug; OEC – oral enzyme combination; TMJ – temporomandibular joint; WOMAC – Western Ontario and McMaster Universities osteoarthritis index.




Fig. 2. Risk-of-bias assessment of the randomized controlled trials (RCTs) included in the review using the Cochrane risk-of-bias (RoB) 2.0 tool

The non-randomized study²³ was evaluated as having a moderate risk of bias, primarily due to limitations regarding confounding factors, participant selection and classification of interventions. Other domains, including deviations from the intended interventions and missing outcome data, were assessed as having a low risk of bias. Overall, the study reflected a moderate risk of bias, indicating methodological limitations associated with its design (Fig. 3).




Fig. 3. Risk-of-bias assessment of the non-randomized study included in the review using the ROBINS-I (Risk Of Bias In Non-randomized Studies of Interventions)

Discussion

Inflammatory conditions such as OA are usually treated with NSAIDs.^{27–29} Although widely used, long-term NSAID therapy is associated with adverse effects.³⁰ The evidence suggests that the use of NSAIDs may be linked to the progression of radiographic knee OA.³¹ Moreover, NSAIDs can interfere with the natural mechanisms of inflammation resolution, such as tissue healing by inhibiting cyclooxygenase-2 activity, which subsequently suppresses

the production of specialized pro-resolving mediators.³² Systemic enzyme therapy, also referred to as treatment with oral enzyme combinations (OECs), has also been studied as a complementary approach to managing inflammation. Evidence supporting the effectiveness of OECs in OA comes from 9 small clinical studies involving up to 268 patients and a systematic review incorporating a meta-analysis of 6 RCTs comparing OEC therapy with diclofenac in patients with knee OA.³³

The studies included in this review demonstrated the potential efficacy of SET in the management of inflammation, with several papers reporting outcomes comparable to those of diclofenac and additional benefits in specific clinical contexts. Akhtar et al.¹⁶ and Bolten et al.¹⁷ showed that OECs were as effective as diclofenac in managing knee joint health, with the study by Bolten et al.¹⁷ reporting a more favorable safety profile and reduced need for rescue medication. These findings were consistent with those of Klein and Kullich²¹ and Singer et al.,²⁵ who also reported comparable efficacy, although Singer et al.²⁵ uniquely highlighted prolonged benefits of OEC therapy after treatment cessation.

Gupta et al.¹⁸ and Jayachandran and Khobre²⁰ focused on TMD and noted improvements in pain, joint function and other clinical outcomes with combination therapy involving OECs and diclofenac. The outcomes of the studies complemented the broader applications of OECs reported in other investigations. These TMJ-specific findings, however, differed in their clinical focus from studies conducted solely on knee and hip joints. For example, Klein et al.²² reported comparable efficacy of OECs and diclofenac in hip OA, as assessed using functional indices such as the Western Ontario and McMaster Universities osteoarthritis index (WOMAC) and the Lequesne algofunctional index (LAFI).

Herrera reported findings that differed from the overall trend, as non-inferiority of OECs to diclofenac could not be established, potentially due to the inadequate sample size, although both treatments showed comparable functional results.¹⁹ Lapi et al. investigated a different outcome, focusing not on treatment efficacy but on the increased risk of acute kidney injury associated with combinations of NSAIDs and antihypertensive medications.²³ Their findings highlighted safety concerns associated with NSAID use, although they did not directly establish the comparative safety of OECs reported by Bolten et al.¹⁷

Tilwe et al.²⁶ provided further evidence suggesting that OECs were superior to diclofenac in reducing joint tenderness, consistent with the studies by Bolten et al.¹⁷ and Singer et al.²⁵ Roth and Stauder reported that diclofenac was modestly superior on the LAFI, although the difference remained within the predefined non-inferiority margin.²⁴ Overall, the majority of studies reported consistent efficacy and safety outcomes, although some discrepancies were observed, mainly due to differences in focus areas and methodological limitations, highlighting the need for more robust trials across various conditions.

The findings of our review, when compared with those of external studies, revealed both similarities and differences in the efficacy and safety of SET and NSAIDs. Bindu et al. highlighted the widespread use of NSAIDs in the management of pain and inflammation, emphasizing their analgesic and anti-inflammatory benefits alongside significant multiorgan adverse effects,³⁴ which were not comprehensively addressed in our review. Szeto et al. proposed a practical framework for minimizing NSAID-related adverse effects through close patient monitoring and specific prescribing strategies,³⁵ which were beyond the clinical scope of our review.

Henrotin et al. reported that OECs were as effective as NSAIDs in the management of OA, with fewer adverse effects.³⁶ These findings are consistent with those of our review, which indicated comparable efficacy of OECs in terms of pain relief and functional outcomes. Furthermore, Ueberall et al. confirmed the comparable effectiveness of OECs and NSAIDs in reducing pain and improving physical function, with significantly more favorable safety and tolerability profiles for OECs than for NSAIDs.³⁷ These findings were consistent with those of Bolten et al.¹⁷ and other knee OA trials included in our review. Shekhar et al. emphasized the benefits of combination therapy for TMJ osteoarthritis, consistent with our findings suggesting greater effectiveness of OECs combined with NSAIDs compared with NSAIDs alone.³⁸ Our review and the available external studies reached similar conclusions regarding the efficacy and safety of SET. However, external studies addressed a broader range of safety concerns and provided practical recommendations for NSAID use, highlighting the need for further research in this area.

Limitations

This review was limited by heterogeneity in study designs, intervention durations and outcome measures. Several studies included small sample sizes, had limited statistical power and lacked comprehensive long-term follow-up. Considerable variation in SET dosing protocols and formulations further hindered direct comparisons between studies. Some studies were assessed as having a moderate risk of bias, particularly the non-randomized trial, which may have affected the validity of the findings. Lastly, the lack of uniform reporting standards across studies made it difficult to comprehensively synthesize the available data.

Future implications

Based on the available evidence, SET may be considered a potential adjunct to NSAIDs in the treatment of certain inflammatory conditions for better efficacy and safety. Establishing standardized SET dosing protocols and

formulations may improve the consistency of therapeutic outcomes. Large-scale, long-term clinical trials are required to provide more robust evidence of the efficacy and safety of treatment across different patient populations. Furthermore, combination therapies involving SET and NSAIDs should be evaluated to determine optimal treatment regimens and investigate their potential to reduce reliance on conventional anti-inflammatory medications.

Conclusions

Systemic enzyme therapy demonstrated efficacy comparable to that of NSAIDs in several studies, with a potentially more favorable safety profile and a lower incidence of adverse effects. In some studies, combining SET with NSAIDs was associated with greater pain relief and improved functional outcomes compared with NSAID treatment alone. These findings suggest that SET may be a useful therapeutic option for selected inflammatory conditions, particularly when reducing treatment-related adverse effects is a clinical priority. However, the strength of the evidence remains limited by differences in study quality, highlighting the need for additional high-quality clinical trials.

Ethics approval and consent to participate

Not applicable.

Data availability

The analyzed data was obtained from publicly available peer-reviewed studies included in the review.

Consent for publication

Not applicable.

Use of AI and AI-assisted technologies

During the preparation of this manuscript, the authors used DeepL Translator (DeepL SE, Cologne, Germany) for language correction. After its use, the authors thoroughly reviewed, verified and revised all content to ensure accuracy and originality.

ORCID iDs

Ganiga Channaiah Shivakumar  <https://orcid.org/0000-0001-7572-9873>
Nishath Sayed Abdul  <https://orcid.org/0000-0003-4091-3632>
Mahesh Shenoy  <https://orcid.org/0000-0001-5486-4566>
Abdullah Abdulaziz Saeidi  <https://orcid.org/0000-0002-0822-993X>
Vincenzo Ronsivalle  <https://orcid.org/0000-0003-3059-4981>
Hande Uzuncibuk  <https://orcid.org/0000-0001-9265-1772>
Gabriele Cervino  <https://orcid.org/0000-0003-4619-4691>
Maria Maddalena Marrapodi  <https://orcid.org/0000-0002-9494-6942>
Giuseppe Minervini  <https://orcid.org/0000-0002-8309-1272>

References

- Chen HS, Cui Y, Zhou ZH, et al.; ARAMIS Investigators. Dual antiplatelet therapy vs alteplase for patients with minor nondisabling acute ischemic stroke: The ARAMIS randomized clinical trial. *JAMA*. 2023;329(24):2135–2144. doi:10.1001/jama.2023.7827
- Salomon-Zimri S, Pushett A, Russek-Blum N, et al. Combination of ciprofloxacin/celecoxib as a novel therapeutic strategy for ALS. *Amyotroph Lateral Scler Frontotemporal Degener*. 2023;24(3–4):263–271. doi:10.1080/21678421.2022.2119868
- Kappos L, Fox RJ, Burcklen M, et al. Ponesimod compared with teriflunomide in patients with relapsing multiple sclerosis in the active-comparator phase 3 OPTIMUM study: A randomized clinical trial. *JAMA Neurol*. 2021;78(5):558–567. doi:10.1001/jamaneurol.2021.0405
- Watanabe H, Morimoto T, Natsuaki M, et al.; STOPDAPT-2 Investigators. Clopidogrel vs aspirin monotherapy beyond 1 year after percutaneous coronary intervention. *J Am Coll Cardiol*. 2024;83(1):17–31. doi:10.1016/j.jacc.2023.10.013
- Harirforoosh S, Asghar W, Jamali F. Adverse effects of nonsteroidal antiinflammatory drugs: An update of gastrointestinal, cardiovascular and renal complications. *J Pharm Pharm Sci*. 2013;16(5):821–847. doi:10.18433/j3vw2f
- Pavan R, Jain S, Shradddha, Kumar A. Properties and therapeutic application of bromelain: A review. *Biotechnol Res Int*. 2012;2012:976203. doi:10.1155/2012/976203
- Hong SJ, Lee SJ, Suh Y, et al.; T-PASS (Ticagrelor Monotherapy in Patients Treated With New-Generation Drug-Eluting Stents for Acute Coronary Syndrome) Investigators. Stopping aspirin within 1 month after stenting for ticagrelor monotherapy in acute coronary syndrome: The T-PASS randomized noninferiority trial. *Circulation*. 2024;149(8):562–573. doi:10.1161/CIRCULATIONAHA.123.066943
- Nishiyama Y, Kimura K, Otsuka T, et al.; CSPS.com Trial Investigators. Dual antiplatelet therapy with cilostazol for secondary prevention in lacunar stroke: Subanalysis of the CSPS.com trial. *Stroke*. 2023;54(3):697–705. doi:10.1161/STROKEAHA.122.039900
- Sands BE, Peyrin-Biroulet L, Kierkus J, et al. Efficacy and safety of mirikizumab in a randomized phase 2 study of patients with Crohn's disease. *Gastroenterology*. 2022;162(2):495–508. doi:10.1053/j.gastro.2021.10.050
- Han MH, Nam JH, Noh E, Lee EK. Gastrointestinal risk of non-steroidal anti-inflammatory drugs and gastroprotective agents used in the treatment of osteoarthritis in elderly patients: A nationwide retrospective cohort study. *Int J Pharmacol Ther*. 2019;57(11):531–541. doi:10.5414/CP203377
- Zobdeh F, Eremenko II, Akan MA, et al. Pharmacogenetics and pain treatment with a focus on non-steroidal anti-inflammatory drugs (NSAIDs) and antidepressants: A systematic review. *Pharmaceutics*. 2022;14(6):1190. doi:10.3390/pharmaceutics14061190
- Bai S, Guo W, Feng Y, et al. Efficacy and safety of anti-inflammatory agents for the treatment of major depressive disorder: A systematic review and meta-analysis of randomised controlled trials. *J Neurol Neurosurg Psychiatry*. 2020;91(1):21–32. doi:10.1136/jnnp-2019-320912
- Page MJ, Moher D, Bossuyt PM, et al. PRISMA 2020 explanation and elaboration: Updated guidance and exemplars for reporting systematic reviews. *BMJ*. 2021;372:n160. doi:10.1136/bmj.n160
- Igelström E, Campbell M, Craig P, Katikireddi SV. Cochrane's risk of bias tool for non-randomized studies (ROBINS-I) is frequently misapplied: A methodological systematic review. *J Clin Epidemiol*. 2021;140:22–32. doi:10.1016/j.jclinepi.2021.08.022
- Sterne JAC, Savović J, Page MJ, et al. RoB 2: A revised tool for assessing risk of bias in randomised trials. *BMJ*. 2019;366:l4898. doi:10.1136/bmj.l4898
- Akhtar NM, Naseer R, Farooqi AZ, Aziz W, Nazir M. Oral enzyme combination versus diclofenac in the treatment of osteoarthritis of the knee – a double-blind prospective randomized study. *Clin Rheumatol*. 2004;23(5):410–415. doi:10.1007/s10067-004-0902-y
- Bolten WW, Glade MJ, Raum S, Ritz BW. The safety and efficacy of an enzyme combination in managing knee osteoarthritis pain in adults: A randomized, double-blind, placebo-controlled trial. *Arthritis*. 2015;2015:251521. doi:10.1155/2015/251521
- Gupta P, Naik SR, Ashok L, Poornima R, Shetty R. Combination of trypsin, rutoside, bromelain and diclofenac sodium in the management of internal derangement of temporomandibular joint: A randomized clinical trial. *J Indian Acad Oral Med Radiol*. 2020;32(3):216–221. doi:10.4103/jiaomr.jiaomr_45_20
- Herrera EG. Phlogenzym in the treatment of a monoarticular painful gonarthrosis: Efficacy and tolerance 1998. <https://web.archive.org/web/20070207173949/https://www.wobenzym.cz/cdweb/mucos/pittompg.htm>. Accessed June 7, 2025.
- Jayachandran S, Khobre P. Efficacy of bromelain along with trypsin, rutoside trihydrate enzymes and diclofenac sodium combination therapy for the treatment of TMJ osteoarthritis – a randomised clinical trial. *J Clin Diagn Res*. 2017;11(6):ZC09–ZC11. doi:10.7860/JCDR/2017/25771.9964
- Klein G, Kullich W. Short-term treatment of painful osteoarthritis of the knee with oral enzymes. *Clin Drug Investig*. 2000;19(1):15–23. doi:10.2165/00044011-200019010-00003
- Klein G, Kullich W, Schnitker J, Schwann H. Efficacy and tolerance of an oral enzyme combination in painful osteoarthritis of the hip: A double-blind, randomised study comparing oral enzymes with non-steroidal anti-inflammatory drugs. *Clin Exp Rheumatol*. 2006;24(1):25–30. PMID:16539815.
- Lapi F, Azoulay L, Yin H, Nessim SJ, Suissa S. Concurrent use of diuretics, angiotensin converting enzyme inhibitors, and angiotensin receptor blockers with non-steroidal anti-inflammatory drugs and risk of acute kidney injury: Nested case–control study. *BMJ*. 2013;346:e8525. doi:10.1136/bmj.e8525
- Roth SH, Stauder GM. Oral enzyme therapy (Phlogenzym) in osteoarthritis: Long-term comparative study against diclofenac. Poster presented at: 65th Annual Scientific Meeting of the American College of Rheumatology; November 11–15, 2001; San Francisco, CA.
- Singer F, Singer C, Oberleitner H. Phlogenzym® versus diclofenac in the treatment of activated osteoarthritis of the knee. A double-blind prospective randomized study. *Int J Immunother*. 2001;17:135–141.
- Tilwe GH, Beria S, Turakhia NH, Daftary GV, Schiess W. Efficacy and tolerability of oral enzyme therapy as compared to diclofenac in active osteoarthritis of knee joint: An open randomized controlled clinical trial. *J Assoc Physicians India*. 2001;49:617–621. PMID:11584936.
- Robbins MS. Diagnosis and management of headache: A review. *JAMA*. 2021;325(18):1874–1885. doi:10.1001/jama.2021.1640
- Amaechi O, Huffman MM, Featherstone K. Pharmacologic therapy for acute pain. *Am Fam Physician*. 2021;104(1):63–72. PMID:34264611.
- Ribeiro H, Rodrigues I, Napoleão L, et al. Non-steroidal anti-inflammatory drugs (NSAIDs), pain and aging: Adjusting prescription to patient features. *Biomed Pharmacother*. 2022;150:112958. doi:10.1016/j.biopha.2022.112958
- Zappavigna S, Cossu AM, Grimaldi A, et al. Anti-inflammatory drugs as anticancer agents. *Int J Mol Sci*. 2020;21(7):2605. doi:10.3390/ijms21072605
- Bakhshi GD, Rangwala ZS, Chauhan AR. Systemic enzyme therapy in chronic venous disease: A review. *Int J Res Med Sci*. 2021;9(9):2869–2876. doi:10.18203/2320-6012.ijrms20213435
- Karnam S, Donthi MR, Jindal AB, Paul AT. Recent innovations in topical delivery for management of rheumatoid arthritis: A focus on combination drug delivery. *Drug Discov Today*. 2024;29(8):104071. doi:10.1016/j.drudis.2024.104071
- Mercer MA, Davis JL, McKenzie HC. The clinical pharmacology and therapeutic evaluation of non-steroidal anti-inflammatory drugs in adult horses. *Animals (Basel)*. 2023;13(10):1597. doi:10.3390/ani13101597
- Bindu S, Mazumder S, Bandyopadhyay U. Non-steroidal anti-inflammatory drugs (NSAIDs) and organ damage: A current perspective. *Biochem Pharmacol*. 2020;180:114147. doi:10.1016/j.bcp.2020.114147
- Szeto CC, Sugano K, Wang JG, et al. Non-steroidal anti-inflammatory drug (NSAID) therapy in patients with hypertension, cardiovascular, renal or gastrointestinal comorbidities: Joint APAGE/APLAR/APSE/APSH/APSN/PoA recommendations. *Gut*. 2020;69(4):617–629. doi:10.1136/gutjnl-2019-319300

36. Henrotin YE, Michlmayr C, Rau SM, Quirke AM, Bigoni M, Ueberall MA. Combination of enzymes and rutin to manage osteoarthritis symptoms: Lessons from a narrative review of the literature. *Rheumatol Ther.* 2022;9(5):1305–1327. doi:10.1007/s40744-022-00472-7
37. Ueberall MA, Mueller-Schwefe GH, Wigand R, Essner U. Efficacy, tolerability, and safety of an oral enzyme combination vs diclofenac in osteoarthritis of the knee: Results of an individual patient-level pooled reanalysis of data from six randomized controlled trials. *J Pain Res.* 2016;9:941–961. doi:10.2147/JPR.S108563
38. Shekhar A, Maddheshiya N, Adit, Rastogi V, Ramalingam K. Anti-inflammatory role of trypsin, rutoside, and bromelain combination in temporomandibular joint osteoarthritis: A systematic review. *Cureus.* 2024;16(1):e51749. doi:10.7759/cureus.51749