

Genomics in dental implantology: The role of genetic and epigenetic factors in dental implant failure – a narrative review

Shilpa Shrinivas Prabhu^{B–E}, Aparna Narayana^{A,F}, Gayathri Krishnamoorthy^{D–F}

Department of Prosthodontics and Crown & Bridge, Manipal College of Dental Sciences, Manipal Academy of Higher Education (MAHE), India

A – research concept and design; B – collection and/or assembly of data; C – data analysis and interpretation;
D – writing the article; E – critical revision of the article; F – final approval of the article

Dental and Medical Problems, ISSN 1644-387X (print), ISSN 2300-9020 (online)

Dent Med Probl.

Address for correspondence

Aparna Narayana
E-mail: aparna.narayan@manipal.edu

Funding sources

None declared

Conflict of interest

None declared

Acknowledgements

None declared

Received on February 21, 2025

Reviewed on May 13, 2025

Accepted on June 19, 2025

Published online on October 31, 2025

Cite as

Prabhu SS, Narayana A, Krishnamoorthy G. Genomics in dental implantology: The role of genetic and epigenetic factors in dental implant failure – a narrative review [published online as ahead of print on October 31, 2025]. *Dent Med Probl.* doi:10.17219/dmp/207334

DOI

10.17219/dmp/207334

Copyright

Copyright by Author(s)

This is an article distributed under the terms of the Creative Commons Attribution 3.0 Unported License (CC BY 3.0) (<https://creativecommons.org/licenses/by/3.0/>).

Abstract

Dental implants are a widely used solution for tooth replacement, yet implant failures remain a challenge. Genetic predispositions and epigenetic modifications influence osseointegration and peri-implant health. The present review explores genetic mechanisms affecting implant healing and introduces implantogenomics – a personalized approach to implant therapy based on an individual's genetic profile.

A comprehensive review of literature from PubMed®, Scopus, EMBASE, and Web of Science (2008–2024) was conducted using Medical Subject Headings (MeSH) terms such as “genetic markers,” “implantogenomics” and “epigenetics.” After removing duplicates and screening for relevance, a total of 46 studies were included in the analysis.

Key genetic variants in bone metabolism (collagen type 1 alpha 1 (*COL1A1*), runt-related transcription factor 2 (*RUNX2*), vitamin D receptor (*VDR*)), immune response (interleukin-1 (*IL-1*), tumor necrosis factor alpha (*TNF-α*), *IL-6*), and osseointegration-related genes (osteoprotegerin (*OPG*), receptor activator of nuclear factor kappa B (*RANK*), receptor activator of nuclear factor kappa-B ligand (*RANKL*)) were identified as potential contributors to implant failure. Epigenetic modifications, including DNA methylation, histone changes and microRNAs (*miRNAs*), regulate bone remodeling and immune responses, and have an influence on implant integration.

Advances in genomics have paved the way for personalized implant therapy through genetic screening, optimizing outcomes and reducing the number of implant failures. Implantogenomics is aimed at tailoring treatments based on genetic profiles, while epigenetic therapies, such as gene modulation, enhance implant integration. Future research should focus on predictive biomarkers and precision-based strategies to improve implant longevity.

Genetic and epigenetic factors play a crucial role in the success of dental implants. Integrating genomic insights into clinical practice can enhance patient selection, predict implant success and improve treatment outcomes. Further research is necessary to establish predictive biomarkers and targeted interventions.

Keywords: genetic factors, epigenetics, implant failure, dental implant, genetic markers

Highlights

- Although dental implants are the preferred option for tooth replacement, failures may still occur due to impaired osseointegration.
- Host-related factors, especially genetic predispositions, significantly influence implant outcomes, with some individuals at higher risk of failure.
- Advances in omics sciences and high-throughput methods enable the identification of molecular mechanisms influencing osseointegration, enabling early risk assessment.
- The emerging field of implantogenomics applies genetic insights to personalize treatment strategies and enhance long-term implant success.

Introduction

Over the past 30 years, dental implants have evolved and become the preferred treatment option for tooth replacement by dentists and patients. The survival rate of implant-supported restorations has increased from 94.6% to 97.1% over the past 2 decades.¹ Despite this high success rate, there has been a predominant increase in the incidence of dental implant failure. Previous studies have noted a dental implant failure rate ranging from 1% to 19%.² Early implant failures have been attributed to altered wound healing, preventing osseointegration, while late implant failures have been associated with extensive peri-implant bone loss after functional loading.³ Understanding the cause of implant failures is essential for their prevention.

Knowledge regarding the factors influencing implant failure and meticulous observation of the implant after placement are crucial elements in the management of the complications such as inflammation, proliferation and progressive bone loss in and around the implant, compromised aesthetics, prosthesis failure, soft tissue dehiscence, implant fracture, and ultimately, implant failure.⁴ Among all variables contributing to dental implant failure, the host factor has emerged as a contentious risk component.² Many researchers have sought to uncover the association between alleles and/or genotypes of genetic markers and the predisposition to implant failure. Analyzing these genetic elements associated with dental implant loss may offer insights into the factors contributing to the varied patient response to currently available treatment options.⁵

Clinicians are able to assess the risk of complications in patients with a negative host response before any elective surgical procedure, as such a response may lead to implant rejection, wherein the host body fails to integrate with the implant. Thus, it is evident that the incidence of implant failure is higher in a subset of individuals who demonstrate a definitive host characteristic, such as genetic factors that disturb the process of osseointegration.⁶ This phenomenon of a small number of patients concentrating risk for implant loss has been termed “clusterization”⁷

High-throughput methodologies are increasingly employed to gain a comprehensive understanding of cellular processes, enabling faster discoveries in health and disease research.⁸

The present review aims to summarize the genetic mechanisms underlying osseointegration healing and introduce implantogenomics, a concept that applies personalized medicine to tailor implant therapy for individual patients based on their unique genetic profile. The study will focus on the following key aspects: genetic factors influencing the prognosis of implant treatment; diagnostic tools for screening high-risk populations; omics profiling in osseointegration; and personalized dental implant therapy. The review will also emphasize the importance of integrating omics sciences with advanced bioengineering technologies to enhance bone formation and regulate osteogenesis by elucidating the genetic and epigenetic signaling cascades involved in dental implantology.

Material and methods

The included studies predominantly focused on genetic and epigenetic contributors to implant failure, with an emphasis on osseointegration. These comprised original research, clinical studies and relevant reviews that examined molecular mechanisms, gene polymorphisms, epigenetic modifications, and the application of omics technologies within the domain of dental implantology. Prominent electronic databases like PubMed®, Scopus, EMBASE, and Web of Science were used to retrieve articles published in the English language during the 16-year period from 2008 to 2024. The time restriction was implemented to preclude the introduction of inaccurate, questionable or outdated concepts while concurrently facilitating the comprehension of the contemporary perception of genetics and its relevance in implant failure. Studies were excluded if they focused solely on mechanical or prosthetic factors, without addressing genetic aspects, or if they lacked peer-reviewed content, including conference abstracts, editorials and opinion articles. A combination of Medical Subject Headings (MeSH)

terms such as “genetic markers”, “implantogenomics”, “genetic polymorphism”, “genetic factors”, “epigenetics”, “peri-implantitis”, “implant failure”, “dental implant”, “precision medicine”, “epigenetic mechanisms”, “DNA methylation”, “histone modification”, and “omics” were used with the Boolean operators to curate the data. Duplicates and methodologically weak studies were excluded during the screening process.

The total number of articles retrieved from 4 online databases was 158. During the screening phase, 40 articles were identified as duplicate entries and were hence removed from the study. In the eligibility phase, 118 records were reviewed, of which 72 were excluded due to deviation from the intended study objective. Finally, 46 articles were included in the review (Fig. 1).

Results and discussion

Peri-implantitis and periodontitis have shown similar clinicopathological features, involving soft tissue damage, infection and bone loss.⁹ An inflammatory process is a notable problem in dental implant patients, as it spreads rapidly and more profoundly around an implant as compared to natural teeth. Therefore, more emphasis should be placed on genes associated with immune/inflammatory responses to foreign bodies.⁶

Wear of the implant surface results in debris-mediated implant loosening, which is one of the main causes of implant failure. This process is referred to as osteolysis. The particles shed from titanium implants trigger a more robust immune response from macrophages when compared to particles derived from supplementary substances used in implant restoration. The inflammatory and osteolytic process of peri-implantitis is driven by pro-inflammatory cytokines, involving interleukin-1 (*IL-1*) and tumor necrosis factor alpha (*TNF-α*), released by macrophages. The evidence indicates that titanium particles cause inflammation and osseous disintegration only in certain individuals receiving implants, which underscores a critical role of the host factor in implant failure.¹⁰




Fig. 1. Flowchart of the literature search process

Genetic factors influencing the prognosis of implant treatment

Genetic mediators that play a vital role in the immune/inflammatory reaction of the body can be categorized as follows (Fig. 2,3):

- ILs;
- bone morphogenetic proteins (BMPs), TNF, transforming growth factor (TGF) (*TNF-α* and *TGF-β*);
- matrix metalloproteinases (MMPs);
- bone metabolism biomarkers.^{9,11}

Interleukins

The diagnostic markers of implant failure include *IL-1A* –889 C/T (rs1800587), *IL-1B* +3954 C/T (rs1143634), *IL-1RN* +2018 T/C (rs419598), and *TNF-α* –308 G/A (rs1800629) genotyping, in vitro *IL-1β*/*TNF-α* release assays, and lymphocyte transformation tests.⁹ Other prognostic markers of peri-implantitis are cathepsin K, receptor activator of nuclear factor kappa-B ligand (*RANKL*) or osteoprotegerin (*OPG*), but further investigations and large clinical trials are necessary to confirm these findings.¹² The amalgamation of *IL-1* allele 2




Fig. 2. Genetic mediators involved in the immune/inflammatory reaction
BMP – bone morphogenetic protein; HMGB – high-mobility group box; IL – interleukin; *IL-1RN* – gene encoding interleukin-1 receptor antagonist; MMP – matrix metalloproteinase; OAF – osteoclast-activating factor; OPG – osteoprotegerin; *RANKL* – receptor activator of nuclear factor kappa-B ligand; *TNF-α* – tumor necrosis factor alpha.




Fig. 3. Effect of genetic mediators on implant surfaces

(*IL-1A* -889 and *IL-1B* +3954) in patients with inflamed periodontal or peri-implant tissues acts as a detrimental factor that exacerbates tissue destruction.¹³ Table 1 summarizes the impact of different ILs on implant prognosis.

Another non-invasive means of inspecting the host's reaction in periodontal and peri-implant diseases is the analysis of gingival crevicular fluid (GCF) or peri-implant sulcular fluid (PISF).¹⁶

TNF and TGF

Single-nucleotide polymorphisms (SNPs) are minor DNA mutations that influence the process of osseointegration in implants. Inflammatory proteins play a crucial role in both the breakdown of the extracellular matrix (ECM) and the resorption of the alveolar bone.⁵ Tumor necrosis factor alpha is a proinflammatory cytokine. It serves as the primary mediator in the immune response to Gram-negative bacteria, with *TNF-α* levels indicating the bacterial load and the severity of inflammation.¹⁶ Bone morphogenetic proteins belong to the *TGF-β* superfamily and have been demonstrated to promote bone ingrowth, facilitate gap healing and enhance implant fixation in various animal studies.¹⁸ BMP-2 and BMP-7 are termed human osteogenic proteins. Table 2 presents an overview of various polymorphic factors and their influence on implant treatment.

Matrix metalloproteinases and extracellular matrix remodeling mediators

The extracellular reaction may vary depending on the implant surface, with ECM exhibiting different morphological characteristics across various material surfaces.

Matrix metalloproteinases are a family of highly conserved endopeptidases. These ECM macromolecules contribute to cellular development and morphogenesis. They regulate growth factors, activate cell surface receptors and influence adhesion molecules. Matrix metalloproteinases are involved in various physiological processes, including inflammatory cell activity, wound healing, angiogenesis, and bone formation.^{20,21}

The family of MMPs consists of 5 groups:

- collagenases: *MMP-1*, *MMP-8*, *MMP-13*, *MMP-18*;
- gelatinases: *MMP-2*, *MMP-9*;
- stromelysins: *MMP-3*, *MMP-10*, *MMP-11*;
- matrilysins: *MMP-7*, *MMP-26*;
- membrane-type (MT) MMPs:
 - 4 transmembrane MMPs: *MMP-14*; *MMP-15*; *MMP-16*; *MMP-24*,
 - 2 glycosylphosphatidylinositol-anchored MMPs: *MMP-17*; *MMP-25*.

The effects of MMPs on ECM remodeling are outlined in Table 3.

Bone metabolism biomarkers

Bone is a metabolically active tissue that undergoes continuous remodeling. This process is driven by the dynamic interaction of osteoblasts and osteoclasts, regulated by a complex network of molecular biomarkers. Table 4 summarizes the impact of these biomarkers on dental implant treatment.

Epigenetic mechanisms

Epigenetics refers to the study of heritable changes in the phenotype that occur without alterations to the

Table 1. Effects of different interleukins (ILs) on dental implant prognosis

Genotype	Effect
<i>IL-1B</i> -511 2/2 (<i>IL-1B</i> allele 2)	early marginal bone loss ¹²
<i>IL-6</i> and <i>IL-8</i>	higher expression in periodontitis sites and peri-implant inflammation ¹³
<i>IL-10</i>	no association with implant failure ¹⁴
<i>IL-17</i>	increased levels in gingivitis and periodontal disease; further investigation needed in peri-implantitis ¹⁵
gene coding for <i>IL-1ra</i>	multiple implant loss ¹⁶
<i>IL-1RN</i> allele 2	implant loss in a Caucasian population ¹⁷
allele 5	implant failure in a Portuguese Caucasian population ¹⁷

IL-1ra – interleukin-1 receptor antagonist; *IL-1RN* – gene encoding *IL-1ra*.

Table 2. Polymorphic factors and their effects on dental implant treatment

Polymorphism	Effect
<i>TNF-α</i>	increases osteoclast activation upon implant placement ¹⁶
OAF	high concentrations lead to increased bone loss and implant failure ¹⁶
<i>BMP-2</i>	stimulates bone ingrowth, gap healing and implant fixation ¹⁸
<i>BMP-7</i>	collagen solution causes a notable increase in BIC and reverse torque resistance following immediate application to implant sockets before insertion of implants ¹⁹

BIC – bone implant contact; BMP – bone morphogenetic protein; *TNF-α* – tumor necrosis factor alpha; OAF – osteoclast-activating factor.

Table 3. Impact of matrix metalloproteinases (MMPs) on extracellular matrix (ECM) remodeling

Matrix metalloproteinase	Effect
<i>MMP-1</i>	initiates bone resorption²²
<i>MMP-2</i> and <i>MMP-8</i>	- elevated levels in GCF and PISF - associated with bone destruction, cavitation, inflammation, and granulation tissue formation - high concentration seen in ECM of diseased implants²⁰
<i>MMP-9</i>	plays a critical role in the development of inflammatory periapical lesions and ECM degradation during the initiation and progression of apical periodontitis²⁰
<i>MMP-18</i>	- diagnostic biomarker in peri-implant diagnostics - detected in PISF by Western blot analysis and associated with early implant failure²³
<i>MMP-25</i>	- cleaves gelatin, type IV collagen and fibronectin - participates in cellular migration and intrusion of ECM and BM associated with peri-implantitis²¹
<i>MMP-26</i>	cleaves fibrinogen and ECM components such as fibronectin, vitronectin, gelatin, and type IV collagen, contributing to peri-implant inflammation²¹

BM – basement membrane; GCF – gingival crevicular fluid; PISF – peri-implant sulcular fluid.

underlying DNA sequence. These changes involve modifications to the chromatin structure, which in turn regulate gene expression independently of base sequence variations.²⁹ Environmental factors such as toxins, microbes, stress, diet, and hormones can alter epigenetic patterns, thereby influencing gene activity and cell behavior. These modifications regulate gene expression by either promoting or silencing transcription, blocking mRNA formation or causing protein post-translational modification.³⁰ Key epigenetic mechanisms include DNA methylation, histone modifications and regulation by non-coding RNAs like microRNAs (*miRNAs*).³¹ DNA methylation silences gene expression and regulates key bone-related genes. Histone modifications, like acetylation and methylation, also control gene activity. Histone deacetylases (HDACs) have been shown to influence bone health by regulating osteoblasts, osteoclasts and bone mass. MicroRNAs are short non-coding RNAs (18–22 nucleotides) that regulate gene expression by degrading or repressing target mRNAs. Increased levels of *miRNAs* suppress gene expression, while decreased levels enhance it. Several *miRNAs*, such as *miR-23a*, *miR-34c* and *miR-133a*, directly influence bone formation by targeting runt-related transcription factor 2 (*RUNX2*), a key transcription factor in the differentiation of osteoblasts.³² Table 5 provides

an overview of key genetic factors and their respective roles at the molecular and biochemical levels in implant integration and failure.

Screening of patients at high risk of implant failure

Some of the diagnostic interventions for the identification of individuals at high risk of implant failure include (Fig. 4):

1. Genome-wide association study: It examines the link between SNPs and traits, with a particular focus on major diseases, by comparing the DNA of individuals with different phenotypes related to a specific trait or disease. Participants are divided into 2 groups: those with the disease (cases); and similar sample without the disease (controls). This method, known as the phenotype-first approach, identifies SNPs in which one allele appears more frequently in the disease group. When a SNP demonstrates a significant association with a specific condition, it is considered to indicate a genomic region that could impact disease risk.³⁶
2. Transcriptional profiling of osseointegration: It highlights the complexity of bone healing, a process that involves interwoven biological stages such as

Table 4. Impact of bone metabolism biomarkers on dental implant treatment

Biomarker	Effect
High-mobility group chromosomal protein <i>HMGB1</i>	promotes the release of cytokines in periodontitis and peri-implantitis
<i>HMGB2</i>	high levels in PI-PISF defend peri-implant tissues against inflammation²¹
<i>BRINP3</i>	risk factor for the development of peri-implantitis in the absence of chronic periodontitis²⁴
Cathepsin K	- determinant of peri-implant tissue health - increased levels in the crevicular fluid are associated with concomitant increase in peri-implant bone loss²⁵
Vitamin D	vitamin D deficiency critically impairs bone integration around implants²⁶
Calcitonin	improves bone maturation around titanium implants²⁷
<i>RANKL</i> and <i>OPG</i>	- detected at high concentrations at peri-implantitis sites - associated with a risk of alveolar bone loss along the entire implant surface²⁸

BRINP – bone morphogenetic proteins/retinoic acid inducible neural-specific protein; HMGB – high-mobility group box; PI-PISF – pro-inflammatory peri-implant sulcular fluid; *RANKL* – receptor activator of nuclear factor kappa-B ligand; *OPG* – osteoprotegerin.

Table 5. Key genetic factors and their influence on dental implants at the molecular and biochemical level

Genetic factor	Molecular/biochemical effect	Impact on dental implant
<i>IL-1</i> (<i>IL-1A/B</i>) polymorphisms	Increased <i>IL-1</i> production promotes a stronger inflammatory response and activates osteoclasts.	higher risk of peri-implantitis, bone resorption and implant failure ¹²
<i>IL-6</i> variations	Elevated <i>IL-6</i> levels amplify inflammation and stimulate osteoclastogenesis.	increased risk of soft tissue inflammation and peri-implant bone loss ¹³
<i>TNF-α</i> polymorphisms (e.g., <i>G-308A</i>)	Overproduction of <i>TNF-α</i> enhances osteoclast differentiation and bone resorption.	greater susceptibility to peri-implantitis and implant failure ¹⁶
<i>RANK/RANKL/OPG</i> pathway genes	Imbalance in <i>RANKL/OPG</i> expression leads to excessive osteoclast activity.	accelerated bone loss around implants, compromised osseointegration ²⁸
<i>VEGF</i> polymorphisms	Altered <i>VEGF</i> expression affects angiogenesis and healing.	impaired vascularization leading to reduced bone healing and implant integration ³³
<i>MMP</i> gene variations	Increased <i>MMP</i> activity degrades ECM and connective tissue.	tissue breakdown around implants, higher risk of implant instability ^{20,22}
<i>COL1A1</i> gene variations	Abnormal collagen production affects the quality of bone matrix.	compromised bone strength and impaired osseointegration ³⁴
<i>RUNX2</i> regulation by <i>miRNAs</i>	Disrupted <i>RUNX2</i> function impairs osteoblast differentiation and bone formation.	delayed or defective bone healing around implants ³⁵

VEGF – vascular endothelial growth factor; *COL1A1* – collagen type 1 alpha 1; *RUNX2* – runt-related transcription factor 2.

- inflammation, osteogenesis and angiogenesis. A comparison of gene expression profiles associated with wound healing and those observed at the post-implant site elucidates the natural delays between gene expression, protein translation and tissue maturation. The histological data indicates that the selected time points for transcriptional analysis effectively capture key early transcriptional events that are crucial in the process of osseointegration.³⁷
3. Real-time polymerase chain reaction (PCR): Diagnostic qualitative PCR is used for a rapid detection of disease-specific nucleic acids, while quantitative PCR measures both the presence and the quantity of a particular DNA sequence in each sample. Both quantitative PCR and DNA microarrays are cutting-edge techniques for the analysis of gene expression, offering researchers a more profound understanding of molecular processes. This understanding supports the development of advanced therapeutic prosthetics for dental implant treatments and applications in tissue engineering biology.³⁸
 4. RNA sequencing: Next-generation sequencing (NGS) is used to identify and quantify RNA in a biological

sample at a specific point in time, enabling the analysis of the dynamic cellular transcriptome.³⁹

5. Genome-wide screening of implant failure by vitamin D deficiency: Genome-wide microarray analyses of implant osseointegration suggest that the unique micro-environment created by implant placement significantly influences multiple gene expression networks, potentially involving peripheral circadian rhythm pathways. Notable interactions between the *NPAS2* gene and cartilage matrix genes have led to a proposed model in which bone marrow mesenchymal cells, through a circadian rhythm-related mechanism, initiate ectopic synthesis of cartilage matrix molecules such as type X collagen without forming actual cartilage tissue at the implant site. Additionally, vitamin D deficiency disrupts these processes, impairing effective bone formation and implant integration.⁴⁰
6. Epigenetic methylation assays: These techniques are used to examine changes in DNA methylation, an important regulatory mechanism that alters gene expression without modifying the underlying DNA sequence. Epigenetic methylation assays facilitate the assessment of the impact of environmental and external factors on gene activity, which contributes to diverse biological processes and disease development.⁴¹

The presented molecular techniques are not currently part of routine clinical practice in implant dentistry. However, they are emerging tools with the potential for future applications as the field of precision diagnostics continues to evolve.

Clinically available tests for genetic screening related to dental implants or periodontal diseases include:

1. MyPerioID® IL-6: saliva-based genetic screening tool that identifies variations of the *IL-6* gene, a critical marker of inflammation. By detecting these genetic predispositions, the test enables the assessment of an individual's risk for developing severe periodontal disease⁴²;




Fig. 4. Diagnostic tools for screening high-risk populations
PCR – polymerase chain reaction.

2. **TNF- α test:** *TNF- α* (G-308A) gene polymorphism has been investigated for its potential association with implant failure and peri-implantitis, as it may influence inflammatory responses. However, research findings have been inconsistent, with some studies showing a possible link and others finding no significant association. Genetic testing for TNF- α variations has yet to be incorporated into standard dental implant planning, as clinical factors like patient health, oral hygiene, bone quality, and surgical technique remain the primary predictors of implant success⁴³;
3. **IL-6 test:** measures salivary IL-6 levels, serving as a biomarker for periodontitis, a major risk factor for dental implant failure. Elevated IL-6 levels indicate increased inflammation and help identify individuals with a higher risk for periodontitis or implant complications. The protocol entails the collection of a saliva sample, the analysis of IL-6 concentrations, and the assessment of the inflammatory status based on the obtained results. Although it is not a genetic test, the IL-6 test provides valuable insight into a patient's risk for implant failure due to inflammation and bone loss⁴⁴;
4. **GenoType Periodontal Susceptibility Test (PST):** a screening tool for *IL-1A* and *IL-1B* gene variations that have been linked to an increased risk of severe periodontitis. A positive result (PST+) indicates higher susceptibility to periodontal disease. While useful for assessing periodontal risk, it is not routinely used in dental implant planning, and evidence supporting its role in predicting peri-implantitis remains limited.⁴⁵
5. **PerioPredict:** genetic risk assessment tool that is used to evaluate moderate to severe periodontal disease by analyzing *IL-1* gene variations, a key factor in inflammation. While it may offer insights for implant planning, it is not specific to dental implants and has shown mixed results in clinical studies. The clinical examination remains the primary method for assessing the periodontal risk. PerioPredict should be used as a complementary tool in conjunction with other clinical evaluations during treatment planning.⁴⁶

Building upon the use of clinically available genetic screening tests, omics profiling is now being explored to gain deeper insights into the mechanisms underlying osseointegration.

Omics profiling in osseointegration

Omics is a branch of biology that encompasses fields like genomics, proteomics, transcriptomics, and metabolomics. The primary goal of omics sciences is to identify, characterize and quantify the diverse biological molecules that contribute to the structure, function and dynamics of cells, tissues or organisms.⁴⁷

Omics technologies have been utilized in distinct pre-clinical studies to understand the early and late molecular events occurring during osseous formation.⁴⁸ Identifying

the genes and proteins that affect osseointegration is essential in order to reduce the healing time associated with implant surgery and improve clinical outcomes in patients with local or systemic conditions that impair bone metabolism.⁴⁹

Studies have shown that an early stage of osseous wound healing was associated with enhanced chemokine, NF- κ B, *TNF- α* signaling pathway, and angiogenesis-related pathways. In the latter stages, an increased expression of mitogen-activated protein kinase (MAPK), Wnt pathways and proteins associated with ECM remodeling and bone mineralization was observed.^{50–53}

A few in vitro studies were conducted to evaluate the impact of different implant surfaces on osteogenic markers, thereby influencing the rate of osseointegration. Moderately rough surfaces exhibited elevated levels of osteogenic markers when compared to polished surfaces.⁵⁴ The hydrophilicity of the implant surface further amplified osteogenesis by positively modulating osteogenesis- and angiogenesis-related pathways (e.g., vascular endothelial growth factor (*VEGF*), MAPK and BMP pathways).⁵⁵

Personalized dental implant therapy

The integration of regenerative approaches to customize the implant therapy as per the patient's individual needs has given rise to a new concept termed implantogenomics or implantomics.⁴⁷

Considerable efforts have been made to create bioactive surface coatings that emulate the biochemical composition and structural characteristics of human bone at the nanoscale. Taking insights from recent omics research, new experimental coatings are currently under development. These coatings are engineered to incorporate targeted drugs, agents, proteins, and growth factors that enhance implant stability by supporting the natural process of osseointegration.⁵⁶ Preclinical studies have shown that coating dental implants with ECM proteins can enhance peri-implant bone formation. De Barros et al. reported increased bone volume and mineralization with collagen type II/chondroitin sulfate coatings in a canine model.⁵⁷ Meng et al. reviewed 34 studies on biomolecular coatings for titanium dental implants, mostly in animal models, and found that growth factors, peptides and ECM proteins may support early stages of bone integration.⁵⁸ However, the authors noted inconsistent results and highlighted the need for clinical studies in humans.⁵⁸ Hasani-Sadrabadi et al. developed a layer-by-layer surface treatment for titanium implants incorporating *BMP-2*-mimicking peptides and gentamicin to enhance osseointegration and antibacterial activity.⁵⁹ Using a poly-dopamine coating to support nanolayer formation, the modified surfaces enabled sustained release of bioactive agents. In vitro and in vivo studies have demonstrated improved cytocompatibility, osteogenic differentiation and

peri-implant bone integration, suggesting promising applications in the fields of dental and orthopedic implantology.⁵⁹ Zhou et al. developed a 3,4-dihydroxyphenylalanine (DOPA)-based peptide coating (DOPA-P1@P2) for titanium implants to address aseptic loosening by promoting staged bone regeneration.⁶⁰ The coating sequentially modulated inflammation, angiogenesis and osteogenesis through specific bioactive peptides. In vivo, it significantly improved push-out strength, bone volume and bone-to-implant contact compared to TiO₂ controls, suggesting its strong potential for enhancing implant osseointegration.⁶⁰

Due to ethical restrictions and an increased prevalence of implant failure, there has been a paucity of in vivo studies investigating the constituents of the host genetic susceptibility that influences biological complications in implant placement.⁴⁹ However, omics technology, which offers a comprehensive understanding of biomaterials, marks a major advance in biomedical science, which will significantly advance the growth of tailored and personalized medicine in implant dentistry.⁶¹ Genetic screening offers potential for personalized dental implant therapy, as it enables the customization of implant design and treatment plans based on a patient's genetic profile. However, challenges such as the complexity of genetic data interpretation, the high cost of testing, and the need for further research to confirm the clinical relevance of genetic markers limit its routine use.

Conclusions

Implant failure is a serious concern in the prognosis of dental implants. Even though the mechanisms underlying implant loss are well-defined, they vary depending on a case. Determining the underlying direct or indirect cause of implant failure is of the utmost importance.

A single-nucleotide polymorphism of pro-inflammatory mediator genes might influence their expression intensity or amino acid sequence, thereby affecting the host inflammatory response. Some SNPs have been correlated with implant loss and determined as probable genetic risk factors for implant failure. Studies on the subject have contributed to the redefinition of prospective targets for successful screening, prevention and maintenance of dental implants. The application of insights from omics sciences has the potential to drive further advancements in personalized dental implant therapy, promoting long-term clinical success.

Ethics approval and consent to participate

Not applicable.

Data availability

Not applicable.

Consent for publication

Not applicable.

Use of AI and AI-assisted technologies

Not applicable.

ORCID iDs

Shilpa Shrinivas Prabhu  <https://orcid.org/0000-0002-8412-9500>
Aparna Narayana  <https://orcid.org/0000-0002-3190-2379>
Gayathri Krishnamoorthy  <https://orcid.org/0000-0001-6816-0754>

References

- Roy M, Loutan L, Garavaglia G, Hashim D. Removal of osseointegrated dental implants: A systematic review of explantation techniques. *Clin Oral Investig*. 2020;24(1):47–60. doi:10.1007/s00784-019-03127-0
- Do TA, Le HS, Shen YW, Huang HL, Fuh LJ. Risk factors related to late failure of dental implant – a systematic review of recent studies. *Int J Environ Res Public Health*. 2020;17(11):3931. doi:10.3390/ijerph17113931
- Oh SL, Shiau HJ, Reynolds MA. Survival of dental implants at sites after implant failure: A systematic review. *J Prosthet Dent*. 2020;123(1):54–60. doi:10.1016/j.prosdent.2018.11.007
- Gupta A, Kale B, Masurkar D, Jaiswal P. Etiology of dental implant complication and failure – an overview. *AIMS Bioengineering*. 2023;10(2):141–152. doi:10.3934/bioeng.2023010
- Cardoso JM, Ribeiro AC, Palos C, Proença L, Noronha S, Alves RC. Association between *IL-1A* and *IL-1B* gene polymorphisms with peri-implantitis in a Portuguese population – a pilot study. *PeerJ*. 2022;10:e13729. doi:10.7717/peerj.13729
- Liao J, Li C, Wang Y, et al. Meta-analysis of the association between common interleukin-1 polymorphisms and dental implant failure. *Mol Biol Rep*. 2014;41(5):2789–2798. doi:10.1007/s11033-014-3133-6
- de Araujo Munhoz FB, Branco FP, Souza RLR, Dos Santos MCLG. Matrix metalloproteinases gene polymorphism haplotype is a risk factor to implant loss: A case-control study. *Clin Implant Dent Relat Res*. 2018;20(6):1003–1008. doi:10.1111/cid.12671
- Attene-Ramos MS, Austin CP, Xia M. High throughput screening. In: Wexler P, ed. *Encyclopedia of Toxicology*. Academic Press; 2014:916–917. doi:10.1016/b978-0-12-386454-3.00209-8
- Zhang F, Finkelstein J. The relationship between single nucleotide polymorphisms and dental implant loss: A scoping review. *Clin Cosmet Investig Dent*. 2019;11:131–141. doi:10.2147/ccide.s207445
- Jacobi-Gresser E, Huesker K, Schütt S. Genetic and immunological markers predict titanium implant failure: A retrospective study. *Int J Oral Maxillofac Surg*. 2013;42(4):537–543. doi:10.1016/j.ijom.2012.07.018
- Ribeiro R, Melo R, Neto PT, Vajgel A, Souza PRE, Címões R. Polymorphisms of *IL-10* (-1082) and *RANKL* (-438) genes and the failure of dental implants. *Int J Dent*. 2017;2017:3901368. doi:10.1155/2017/3901368
- Vaz P, Gallas MM, Braga AC, Sampaio-Fernandes JC, Felino A, Tavares P. *IL1* gene polymorphisms and unsuccessful dental implants. *Clin Oral Implants Res*. 2011;23(12):1404–1413. doi:10.1111/j.1600-0501.2011.02322.x
- Hamdy AAEM, Ebrahim MAE. The effect of interleukin-1 allele 2 genotype (*IL-1a*(-889) and *IL-1b*(+3954)) on the individual's susceptibility to peri-implantitis: Case-control study. *J Oral Implantol*. 2010;37(3):325–334. doi:10.1563/aaid-joi-d-09-00117.1
- Severino VO, Napimoga MH, de Lima Pereira SA. Expression of *IL-6*, *IL-10*, *IL-17* and *IL-8* in the peri-implant crevicular fluid of patients with peri-implantitis. *Arch Oral Biol*. 2011;56(8):823–828. doi:10.1016/j.archoralbio.2011.01.006
- Pigossi SC, Alvim-Pereira F, Montes CC, et al. Genetic association study between interleukin 10 gene and dental implant loss. *Arch Oral Biol*. 2012;57(9):1256–1263. doi:10.1016/j.archoralbio.2012.02.020

16. Petković AB, Matić SM, Stamatović NV, et al. Proinflammatory cytokines (IL-1 β and TNF- α) and chemokines (IL-8 and MIP-1 α) as markers of peri-implant tissue condition. *Int J Oral Maxillofac Surg.* 2010;39(5):478–485. doi:10.1016/j.ijom.2010.01.014
17. Montes CC, Alvim-Pereira F, de Castilhos BB, Sakurai MLL, Olandoski M, Trevilatto PC. Analysis of the association of IL1B(C+3954T) and IL1RN(intron 2) polymorphisms with dental implant loss in a Brazilian population. *Clin Oral Implants Res.* 2009;20(2):208–217. doi:10.1111/j.1600-0501.2008.01629.x
18. Thorey F, Menzel H, Lorenz C, Gross G, Hoffmann A, Windhagen H. Osseointegration by bone morphogenetic protein-2 and transforming growth factor beta2 coated titanium implants in femora of New Zealand white rabbits. *Indian J Orthop.* 2011;45(1):57–62. doi:10.4103/0019-5413.73659
19. Tonetti MS, Hämmerle CHF; European Workshop on Periodontology Group C. Advances in bone augmentation to enable dental implant placement: Consensus Report of the Sixth European Workshop on Periodontology. *J Clin Periodontol.* 2008;35(8 Suppl):168–172. doi:10.1111/j.1600-051x.2008.01268.x
20. Luchian I, Goriuc A, Sandu D, Covasa M. The role of matrix metalloproteinases (MMP-8, MMP-9, MMP-13) in periodontal and peri-implant pathological processes. *Int J Mol Sci.* 2022;23(3):1806. doi:10.3390/ijms23031806
21. Kuula H, Salo T, Pirilä E, et al. Human β -defensin-1 and -2 and matrix metalloproteinase-25 and -26 expression in chronic and aggressive periodontitis and in peri-implantitis. *Arch Oral Biol.* 2008;53(2):175–186. doi:10.1016/j.archoralbio.2007.09.010
22. Leite MFF, Santos MCLG, de Souza AP, Line SRP. Osseointegrated implant failure associated with MMP-1 promoter polymorphisms (-1607 and -519). *Int J Oral Maxillofac Implants.* 2008;23(4):653–658. PMID:18807561.
23. Kim KA, Chung SB, Hawng EY, et al. Correlation of expression and activity of matrix metalloproteinase-9 and -2 in human gingival cells of periodontitis patients. *J Periodontol Implant Sci.* 2013;43(1):24–29. doi:10.5051/jpis.2013.43.1.24
24. Luo L, Xie P, Gong P, Tang X, Ding Y, Deng LX. Expression of HMGB1 and HMGN2 in gingival tissues, GCF and PICF of periodontitis patients and peri-implantitis. *Arch Oral Biol.* 2011;56(10):1106–1111. doi:10.1016/j.archoralbio.2011.03.020
25. Casado PL, Aguiar DP, Costa LC, et al. Different contribution of BRINP3 gene in chronic periodontitis and peri-implantitis: A cross-sectional study. *BMC Oral Health.* 2015;15:33. doi:10.1186/s12903-015-0018-6
26. Dirschnabel AJ, Alvim-Pereira F, Alvim-Pereira CC, Bernardino JF, Rosa EAR, Trevilatto PC. Analysis of the association of IL1B(C-511T) polymorphism with dental implant loss and the clusterization phenomenon. *Clin Oral Implants Res.* 2011;22(11):1235–1241. doi:10.1111/j.1600-0501.2010.02080.x
27. Chen BL, Xie DH, Zheng ZM, et al. Comparison of the effects of alendronate sodium and calcitonin on bone-prosthesis osseointegration in osteoporotic rats. *Osteoporosis Int.* 2010;22(1):265–270. doi:10.1007/s00198-010-1186-5
28. Chaparro A, Beltrán V, Betancur D, et al. Molecular biomarkers in peri-implant health and disease: A cross-sectional pilot study. *Int J Mol Sci.* 2022;23(17):9802. doi:10.3390/ijms23179802
29. Martos SN, Tang WY, Wang Z. Elusive inheritance: Transgenerational effects and epigenetic inheritance in human environmental disease. *Prog Biophys Mol Biol.* 2015;118(1–2):44–54. doi:10.1016/j.pbiomolbio.2015.02.011
30. Williams SD, Hughes TE, Adler CJ, Brook AH, Townsend GC. Epigenetics: A new frontier in dentistry. *Aust Dent J.* 2014;59 Suppl 1:23–33. doi:10.1111/adj.12155
31. Jones PA, Liang G. Rethinking how DNA methylation patterns are maintained. *Nat Rev Genet.* 2009;10(11):805–811. doi:10.1038/nrg2651
32. Di Gianfilippo R, Di Gianfilippo C, Prato GPP. The role of epigenetics on dental implant therapy: A systematic review. *Epigenomes.* 2017;1(2):12. doi:10.3390/epigenomes1020012
33. Palermo A, Ferrante F, Stanca E, et al. Release of VEGF from dental implant surface (IML® Implant) coated with concentrated growth factors (CGF) and the liquid phase of CGF (LPCGF): In vitro results and future expectations. *Appl Sci.* 2019;9(10):2114. doi:10.3390/app9102114
34. Roy M, Corti A, Daniele S, et al. Early changes of ECM-related gene expression in fibroblasts cultured on TiO₂, ZrO₂ and PEEK: The beneficial effects of UVC photofunctionalization. *J Photochem Photobiol.* 2021;6:100023. doi:10.1016/j.jpap.2021.100023
35. Liu F, Wang Z, Liu F, et al. MicroRNA-29a-3p enhances dental implant osseointegration of hyperlipidemic rats via suppressing dishevelled 2 and frizzled 4. *Cell Biosci.* 2018;8:55. doi:10.1186/s13578-018-0254-y
36. Welter D, MacArthur J, Morales J, et al. The NHGRI GWAS Catalog, a curated resource of SNP-trait associations. *Nucleic Acids Res.* 2013;42(D1):D1001–D1006. doi:10.1093/nar/gkt1229
37. Ivanovski S, Hamlet S, Salvi GE, et al. Transcriptional profiling of osseointegration in humans. *Clin Oral Implants Res.* 2011;22(4):373–381. doi:10.1111/j.1600-0501.2010.02112.x
38. Valones MAA, Guimarães RL, Brandão LAC, de Souza PRE, Carvalho AAT, Crovela S. Principles and applications of polymerase chain reaction in medical diagnostic fields: A review. *Braz J Microbiol.* 2009;40(1):1–11. doi:10.1590/s1517-83822009000100001
39. Wang Z, Gerstein M, Snyder M. RNA-Seq: A revolutionary tool for transcriptomics. *Nat Rev Genet.* 2009;10(1):57–63. doi:10.1038/nrg2484
40. Mengatto CM, Mussano F, Honda Y, Colwell CS, Nishimura I. Circadian rhythm and cartilage extracellular matrix genes in osseointegration: A genome-wide screening of implant failure by vitamin D deficiency. *PLoS One.* 2011;6(1):e15848. doi:10.1371/journal.pone.0015848
41. Yong WS, Hsu FM, Chen PY. Profiling genome-wide DNA methylation. *Epigenetics Chromatin.* 2016;9:26. doi:10.1186/s13072-016-0075-3
42. Srivastava N, Nayak PA, Rana S. Point of care – a novel approach to periodontal diagnosis – a review. *J Clin Diagn Res.* 2017;11(8):ZE01–ZE06. doi:10.7860/jcdr/2017/26626.10411
43. Bono A, Brunotto MN, Ibanez JC, et al. Failure of dental implants and its association with IL-10 A/G rs1800896 and TNF- α G308A rs1800629308 polymorphisms, systematic review, and meta-analysis of clinical trials. *Eur J Dent Oral Health.* 2025;6(2):1–7. doi:10.24018/ejdent.2025.6.2.370
44. Drafta S, Guita DM, Cristache CM, et al. Could pro-inflammatory cytokines levels IL-6, IL-8, TNF α , total antioxidant status and lactate dehydrogenase be associated with peri-implant bone loss? A pilot study. *Appl Sci.* 2021;11(22):11012. doi:10.3390/app112211012
45. Brodzikowska A, Górska R, Kowalski J. Interleukin-1 genotype in periodontitis. *Arch Immunol Ther Ex.* 2019;67(6):367–373. doi:10.1007/s00005-019-00555-4
46. Diehl SR, Kuo F, Hart TC. Interleukin 1 genetic tests provide no support for reduction of preventive dental care. *J Am Dent Assoc.* 2015;146(3):164–173.e4. doi:10.1016/j.adaj.2014.12.018
47. Refai AK, Cochran DL. Harnessing omics sciences and biotechnologies in understanding osseointegration— personalized dental implant therapy. *Int J Oral Maxillofac Implants.* 2020;35(3):e27–e39. doi:10.11607/jomi.7272
48. Donos N, Dereka X, Mardas N. Experimental models for guided bone regeneration in healthy and medically compromised conditions. *Periodontol 2000.* 2015;68(1):99–121. doi:10.1111/prd.12077
49. Calciolari E, Donos N. The use of omics profiling to improve outcomes of bone regeneration and osseointegration. How far are we from personalized medicine in dentistry? *J Proteomics.* 2018;188:85–96. doi:10.1016/j.jpro.2018.01.017
50. Wise JK, Sena K, Vranizan K, et al. Temporal gene expression profiling during rat femoral marrow ablation-induced intramembranous bone regeneration. *PLoS One.* 2010;5(10):e12987. doi:10.1371/journal.pone.0012987
51. Ivanovski S, Hamlet S, Retzepi M, Wall I, Donos N. Transcriptional profiling of “guided bone regeneration” in a critical-size calvarial defect. *Clin Oral Implants Res.* 2011;22(4):382–389. doi:10.1111/j.1600-0501.2010.02104.x
52. Donos N, Retzepi M, Wall I, Hamlet S, Ivanovski S. In vivo gene expression profile of guided bone regeneration associated with a microrough titanium surface. *Clin Oral Implants Res.* 2011;22(4):390–398. doi:10.1111/j.1600-0501.2010.02105.x
53. Al-Kattan R, Retzepi M, Calciolari E, Donos N. Microarray gene expression during early healing of GBR-treated calvarial critical size defects. *Clin Oral Implants Res.* 2016;28(10):1248–1257. doi:10.1111/clr.12949

54. Ekdahl KN, Hong J, Hamad OA, Larsson R, Nilsson B. Evaluation of the blood compatibility of materials, cells, and tissues: Basic concepts, test models, and practical guidelines. *Adv Exp Med Biol.* 2012;735:257–270. doi:10.1007/978-1-4614-4118-2_18
55. Donos N, Hamlet S, Lang NP, et al. Gene expression profile of osseointegration of a hydrophilic compared with a hydrophobic microrough implant surface. *Clin Oral Implants Res.* 2011;22(4):365–372. doi:10.1111/j.1600-0501.2010.02113.x
56. Smeets R, Stadlinger B, Schwarz F, et al. Impact of dental implant surface modifications on osseointegration. *Biomed Res Int.* 2016;2016:6285620. doi:10.1155/2016/6285620
57. De Barros RRM, Novaes AB, Korn P, et al. Bone formation in a local defect around dental implants coated with extracellular matrix components. *Clin Implant Dent Relat Res.* 2013;17(4):742–757. doi:10.1111/cid.12179
58. Meng HW, Chien EY, Chien HH. Dental implant bioactive surface modifications and their effects on osseointegration: A review. *Biomark Res.* 2016;4:24. doi:10.1186/s40364-016-0078-z
59. Hasani-Sadrabadi MM, Pouraghaei S, Zahedi E, et al. Antibacterial and osteoinductive implant surface using layer-by-layer assembly. *J Dent Res.* 2021;100(10):1161–1168. doi:10.1177/00220345211029185
60. Zhou W, Liu Y, Nie X, et al. Peptide-based inflammation-responsive implant coating sequentially regulates bone regeneration to enhance interfacial osseointegration. *Nat Commun.* 2025;16(1):3283. doi:10.1038/s41467-025-58444-8
61. Cranford SW, de Boer J, van Blitterswijk C, Buehler MJ. Materiomics: An -omics approach to biomaterials research. *Adv Mater.* 2013;25(6):802–824. doi:10.1002/adma.201202553