

Relationship between the *TAS2R38* and *TAS1R2* polymorphisms and the dental status in obese children

Münevver Kiliç^{1,A–F}, Taskin Gurbuz^{1,A–F}, Cigdem Yuce Kahraman^{2,A–F}, Atilla Cayir^{3,A–F}, Abdülbaki Bilgiç^{4,C,E,F}, Yusuf Kurt^{5,B,C}

¹ Department of Pediatric Dentistry, Faculty of Dentistry, Atatürk University, Erzurum, Turkey

² Department of Medical Genetics, Faculty of Medicine, Atatürk University, Erzurum, Turkey

³ Erzurum Regional Education and Research Hospital, Turkey

⁴ Department of Agricultural Economics, College of Agriculture, Atatürk University, Erzurum, Turkey

⁵ Department of Molecular Biology and Genetics, Harran University, Şanlıurfa, Turkey

A – research concept and design; B – collection and/or assembly of data; C – data analysis and interpretation;

D – writing the article; E – critical revision of the article; F – final approval of the article

Dental and Medical Problems, ISSN 1644-387X (print), ISSN 2300-9020 (online)

Dent Med Probl. 2022;59(2):233–240

Address for correspondence

Münevver Kiliç

E-mail: mnvvr_25@hotmail.com

Funding sources

The Coordination Unit of Scientific Research Projects at Atatürk University, Erzurum, Turkey.

Conflict of interest

None declared

Acknowledgements

The co-authors thank Prof. Yusuf Kurt and Prof. Abdülbaki Bilgiç for statistical assistance.

Received on June 6, 2021

Reviewed on October 15, 2021

Accepted on October 21, 2021

Published online on May 4, 2022

Cite as

Kiliç M, Gurbuz T, Kahraman CY, Cayir A, Bilgiç A, Kurt Y. Relationship between the *TAS2R38* and *TAS1R2* polymorphisms and the dental status in obese children. *Dent Med Probl.* 2022;59(2):233–240. doi:10.17219/dmp/143252

DOI

10.17219/dmp/143252

Copyright

© 2022 by Wrocław Medical University

This is an article distributed under the terms of the

Creative Commons Attribution 3.0 Unported License (CC BY 3.0)

(<https://creativecommons.org/licenses/by/3.0/>).

Abstract

Background. The sweet taste and bitter taste genes are thought to have an influence on obesity and caries, which are chronic diseases.

Objectives. The aim of the study was to investigate the effects of the polymorphisms of *TAS2R38* (the bitter taste gene) and *TAS1R2* (the sweet taste gene), which are the most important members of the taste gene family, on the dental status of obese and normal-weight children.

Material and methods. The study included 78 healthy children and 100 children diagnosed with obesity (5–16 years old). The anthropometric measurements and dental status of the children were evaluated. The decayed, missing and filled permanent/primary teeth (DMFT/dmft) index was determined using the standard methods recommended by the World Health Organization (WHO). Blood samples were collected from all subjects and were analyzed via the polymerase chain reaction (PCR) test, with the use of specific primers for the genetic analysis. Five single-nucleotide polymorphisms (SNPs) of the *TAS2R38* and *TAS1R2* genes were investigated. The truncated Poisson and truncated negative binomial modeling approaches were used with regard to the data.

Results. The DMFT/dmft scores were low in obese children and high in children who did not sense the bitter taste (non-tasters). While obese non-taster children had increased DMFT/dmft scores, normal-weight non-taster children had decreased DMFT/dmft scores.

Conclusions. The alanine, valine and isoleucine (AVI) as well as proline, alanine and valine (PAV) haplotypes of the *TAS2R38* gene are associated with the DMFT/dmft index and obesity. This study showed that the DMFT/dmft scores were decreased in obese children. According to the haplotype analysis of the *TAS2R38* gene, the DMFT/dmft scores were increased in non-tasters. When differentiating obese non-tasters and control non-tasters, DMFT/dmft increased in obese non-taster patients, while it decreased in control non-taster patients.

Keywords: genetic, childhood obesity, tooth decay, oral health, *TAS2R38* polymorphism

Introduction

Caries and obesity are multifactorial diseases having a genetic, epigenetic and dietetic background.¹ Like tooth decay, obesity has become an almost global epidemic among children.² Consuming food with a high carbohydrate (CH) content is closely associated with obesity. Therefore, the CH-weighted eating habits represent a common problem related to tooth decay and obesity.³ There is an apparent correlation between caries and CH consumption as a severe risk factor for obesity.⁴

Carbohydrate consumption, which actively contributes to caries and obesity, may be affected by the genetic code determining our taste perception.⁵ The sense of taste helps humans predict whether the food they consume is beneficial or harmful. The taste has developed as a guide to identify the nutrients needed by the body.⁶ Humans can be categorized into 2 groups with regard to their reactions to the bitter taste – tasters and non-tasters. The most critical component responsible for the bitter taste is phenylthiocarbamide (PTC).⁷ Individuals sensitive to PTC are also susceptible to more than 10,000 other tastes, which is not observed in individuals who are not sensitive to PTC.⁸ The primary gene responsible for PTC sensitivity is *TAS2R38*, located on chromosome 7.⁹ The polymorphisms of the *TAS2R38* bitter taste gene may lead to differences in how the taste of food is sensed. Therefore, any polymorphism of this gene may affect nutritional habits, which play significant roles in the development of tooth decay and obesity.¹⁰ Studies are being carried out to discover the effect of *TAS2R38* on the body mass index (BMI).¹¹ In a recent study, it was shown that the daily energy intake of the ‘non-taster’ group was higher in pre-adolescents as compared to the ‘taster’ group, which enjoys the PTC/6-n-propylthiouracil (PROP) taste.¹² It indicates that *TAS2R38* can be associated with increased BMI and obesity risk in children.¹³ Although the *TAS2R38* polymorphisms have been studied in different populations, there is a lack of data with regard to the Turkish population.

Tooth decay, which is caused by many factors, is one of the most common chronic diseases of childhood. Tooth decay occurs as a result of the complex interaction of many environmental and genetic factors.¹⁴ The relationship between *TAS2R38*, obesity and caries has been stated in the literature.¹⁵ Studies on *TAS2R38* show that low taste sensitivity is associated with a greater risk of caries as compared to high taste sensitivity, which suggests that *TAS2R38* may be associated with caries.¹⁶ The *TAS1R* and *TAS2R* gene families encoding taste receptors are involved in the molecular mechanism of taste perception. The *TAS1R* gene family includes the genes encoding receptors that allow the perception of sweet taste; it comprises 3 taste receptor genes on chromosome 1 – *TAS1R1*, *TAS1R2* and *TAS1R3*.¹⁷ The perception of the sweet taste is mediated by the heterodimeric G protein-coupled receptors encoded by *TAS1R2* and *TAS1R3*.¹⁷ A significant relationship between *TAS1R2*

and caries was observed in a recent study.¹⁸ Yet another study, conducted on twins, showed that genetic differences between individuals influenced taste perception and nutritional habits.¹⁹ However, the research on the topic is still scarce in the literature.

The aim of the present study was to investigate the effects of obesity and the polymorphisms of the *TAS2R38* and *TAS1R2* genes, the major members of the taste gene family, on tooth decay in children through evaluating the caries levels in obese and non-taster children. To this end, 2 hypotheses were tested:

- hypothesis I: There is no difference in the incidence of tooth decay between children with obesity and those of normal weight; and
- hypothesis II: There is no difference in the incidence of tooth decay based on the occurrence of the *TAS2R38* bitter taste and *TAS1R2* sweet taste gene polymorphisms.

Material and methods

This cross-sectional, observational study was approved by the Ethics Committee of the Health Sciences Institute at Atatürk University, Erzurum, Turkey (No. B.30.2.ATA.0A10000-617), and was supported by the Scientific Research Projects Commission at Atatürk University (project No. 2012/66 and 2012/375). The study was carried out in collaboration with the Department of Pediatric Endocrinology and the Department of Medical Genetics of the Faculty of Medicine, and the Department of Pediatric Dentistry of the Faculty of Dentistry at Atatürk University. All participants and their parents were informed about the study aims and procedures before any interventions were made, and written consent was obtained from all parents to confirm their approval for treatment. A patient follow-up form was completed for each participant, and the oral and general health status, weight and height measurements, regular medication use, previous dental treatment, and the eating habits over the previous 3 days were recorded. A total of 100 patients aged 5–16 years who had been diagnosed with obesity after referral to the Department of Pediatric Endocrinology were transferred to the pedodontist liable for the assessment of their oral care (MK). The blood samples (2 mL) obtained from the patients as part of routine examinations were collected into ethylenediaminetetraacetic acid (EDTA)-containing tubes and transferred to the Department of Medical Genetics for the genetic analysis. The control group comprised 78 healthy children aged 6–16 years who had been referred to the Department of Pediatric Dentistry and had no systemic disorders. The same procedures were followed for the control group.

Weight measurements were performed using an electronic scale with a ± 100 g sensitivity range when the participants were wearing lightweight clothes. Each participant stood on the scale with both feet placed at the center

of the platform and the value presented on the electronic screen was recorded in kg.

A steel tape measure was used to determine the height of the children while standing upright with their bare feet parallel and adjacent to each other, and with their shoulders and hips touching the wall. Height measurements were recorded in cm. The growth curves described for Turkish children were used to assess BMI,²⁰ and the participants with BMI corresponding to the 95th percentile or above of age-specific limits were considered obese. The decayed, missing and filled teeth of the participants were identified, and the data was recorded for permanent and primary teeth based on the DMFT/dmft scale, respectively.

DNA isolation and SNP genotyping

Whole blood samples were taken from the children for DNA isolation. The blood samples were added by means of venipuncture into Vacutainer[®] EDTA tubes (BD, Franklin Lakes, USA) and used for manual DNA isolation, utilizing the QIAamp[®] DNA Blood Mini Kit (Qiagen, Hilden, Germany) according to the manufacturer's instructions. A single-nucleotide polymorphism (SNP) identification assay (LightSNiP; TIB Molbiol, Berlin, Germany) employing the simple-probe fluorescence melting curve analysis (FMCA) was used to detect the *TAS2R38* and *TAS1R2* polymorphisms. The Rotor-Gene Q (Qiagen) was used in the polymerase chain reaction (PCR) and high-resolution FMCA procedures. A 20-microliter mixture was prepared for each SNP; it included 1 μ L of LightSNiP reagent mix (TIB Molbiol), 2 μ L of LightCycler[®] FastStart DNA Master HybProbe (Roche Diagnostics, Mannheim, Germany), 1.6 μ L of MgCl₂ (25 mM), and a volume of 50 ng DNA and H₂O. The PCR program included the initial denaturation at 95°C for 10 min, followed by 45 cycles of 10-second denaturation at 95°C, 10-second annealing at 60°C and elongation at 72°C for 15 s. The high-resolution FMCA program included cycles at 95°C for 30 s, at 40°C for 120 s and a ramp to 75°C at a transition rate of 1.5°C/s, with the continuous collection of fluorescence. The amplification and melting curves were generated using the Rotor-Gene Q software. As a result of the analysis, the allelic discrimination of the SNPs was achieved.

Statistical analysis

The obtained data was analyzed with the use of the IBM SPSS Statistics for Windows software, v. 21.0 (IBM Corp., Armonk, USA). Descriptive statistics were calculated for each variable and expressed as mean \pm standard deviation ($M \pm SD$). First, the Kolmogorov–Smirnov and Levene tests were applied to determine whether the data showed normal distribution and the homogeneity of variances. The χ^2 test was used to evaluate gender distribution for children in the obese and control groups. The one-way analysis of variance (ANOVA) was used to evaluate

differences with regard to age, DMFT/dmft and body mass index percentile (BMIP). The abovementioned software package was used to analyze the data, and the results were evaluated at an alpha level of 0.05 within a 95% confidence interval (CI). The Poisson and negative binomial models were applied to the data, using the R program (<https://www.r-project.org>). The DMFT/dmft score was defined as the dependent variable, and obesity and the absence of the bitter taste were defined as the independent variables. Caries was modeled with the obesity and non-taster dummy variables by using the truncated Poisson and truncated negative binomial modeling approaches.

The obesity and non-taster dummy variables were created to denote obesity. Obese and non-taster children were coded as 1, and normal-weight children (control) and tasters were coded as 0. Obese non-tasters and control non-tasters were included in the model and labeled as the obese_non-taster and control_non-taster dummy variables (obese non-tasters and control non-tasters were coded as 1, and obese tasters and control tasters were coded as 0). Considering that the highest number of decayed teeth would be 28 in the study, given the maximum age of 16 years in the study population, both models were truncated in the right truncate. The Wald test was used to assess whether the model coefficients were statistically different from zero.²¹

Results

There were no statistically significant differences in gender distribution ($p = 0.881$) or the mean age ($p = 0.312$ and $p = 0.165$ for boys and girls, respectively) between the obese and control groups. No statistical difference was found between the groups in terms of DMFT/dmft. The mean BMIP value was significantly higher in the obese group than in the control group ($p < 0.001$) (Table 1).

Table 1. Gender distribution, age, the number of decayed, missing and filled permanent/primary teeth (DMFT/dmft), and the body mass index percentile (BMIP) values for obese and control patients

Characteristic		Obese group	Control group	<i>p</i> -value
Gender distribution <i>n</i> (%)	total	100 (56.18)	78 (43.82)	0.881
	M	46 (56.79)	35 (43.21)	
	F	54 (55.67)	43 (44.33)	
Age [years] <i>M</i> \pm <i>SD</i>	total	11.80 \pm 2.53	10.89 \pm 3.50	0.050
	M	11.45 \pm 2.34	10.71 \pm 3.67	0.312
	F	12.16 \pm 2.74	11.05 \pm 3.55	0.165
DMFT/dmft <i>M</i> \pm <i>SD</i>		2.80 \pm 3.13	3.29 \pm 3.93	0.352
BMIP <i>M</i> \pm <i>SD</i>		96.27 \pm 3.79	66.22 \pm 14.36	0.000*

M – male; F – female; *M* – mean; *SD* – standard deviation; *statistically significant.

TAS1R2 genotypes and the DMFT/dmft score

When the genotypes of *rs4920566* SNP were evaluated, the highest DMFT/dmft score was found in the AG heterozygous control group. When the genotypes of *rs9701796* SNP were evaluated, the highest DMFT/dmft value was found in the CC homozygous obese group. However, the difference between the genotypes were not statistically significant (Table 2).

TAS2R38 genotypes and the DMFT/dmft score

When the genotypes of *rs713598* SNP were evaluated, the highest DMFT/dmft score was found in the homozygous control group. However, the difference between the genotypes was not statistically significant. When the genotypes of *rs1726866* SNP and *rs10246939* SNP were evaluated, the highest DMFT/dmft scores were observed in the homozygous control group, but the differences between the genotypes were not statistically significant (Table 3).

TAS2R38 haplotypes and the DMFT/dmft score

When the *TAS2R38* haplotypes were evaluated, no statistical differences were found between the control and obese groups. The highest DMFT/dmft score was found in the CC homozygous genotype in all 3 SNPs of the *TAS2R38* gene (Table 3).

Haplotype frequency

A total of 3 different SNPs of the *TAS2R38* gene were evaluated and the linkage disequilibrium (LD) test was

Table 2. Distribution of the DMFT/dmft scores in the *TAS1R2* genotypes

<i>TAS1R2</i> SNP	Genotype	<i>n</i>	DMFT/dmft score		<i>p</i> -value
			<i>M</i> ± <i>SD</i>	min–max	
<i>rs4920566</i>	AA c	17	3.88 ±3.05	0–10	0.294
	AG c	30	4.10 ±4.80	0–15	
	GG c	31	2.67 ±2.70	0–10	
	AA o	43	2.30 ±2.51	0–8	
	AG o	23	3.09 ±4.69	0–20	
	GG o	34	2.82 ±3.08	0–11	
<i>rs9701796</i>	CC c	2	3.00 ±0.70	2–4	0.354
	CG c	17	2.94 ±3.28	0–10	
	GG c	59	3.04 ±3.92	0–15	
	CC o	3	6.00 ±0.00	6–6	
	CG o	30	3.13 ±4.29	0–20	
	GG o	67	2.39 ±2.74	0–11	

SNP – single-nucleotide polymorphism; min – minimum; max – maximum; c – control group; o – obese group.

Table 3. Distribution of the DMFT/dmft scores in the *TAS2R38* genotypes

<i>TAS2R38</i> SNP	Genotype	<i>n</i>	DMFT/dmft score		<i>p</i> -value
			<i>M</i> ± <i>SD</i>	min–max	
<i>rs713598</i>	CC c	24	4.21 ±3.83	0–20	0.222
	CG c	40	2.98 ±3.54	0–8	
	GG c	14	3.64 ±4.14	0–10	
	CC o	44	3.19 ±4.09	0–15	
	CG o	31	2.39 ±2.45	0–14	
	GG o	25	1.88 ±2.35	0–15	
<i>rs1726866</i>	CC c	20	4.75 ±3.85	0–11	0.270
	CT c	37	2.89 ±3.16	0–15	
	TT c	21	3.29 ±4.39	0–20	
	CC o	27	2.89 ±2.75	0–8	
	CT o	44	2.70 ±3.01	0–11	
	TT o	29	2.38 ±4.04	0–20	
<i>rs10246939</i>	CC c	20	4.45 ±4.05	0–11	0.346
	CT c	37	3.05 ±3.10	0–15	
	TT c	21	3.29 ±4.39	0–20	
	CC o	47	2.32 ±2.94	0–11	
	CT o	23	3.26 ±2.80	0–8	
	TT o	30	2.73 ±4.11	0–20	

performed (Fig. 1) to determine the haplotypes of the respondent children (Table 4). Partial tasters were included in the taster group.




Fig. 1. Linkage disequilibrium (LD) plot for 3 single-nucleotide polymorphisms (SNPs) of *TAS2R38* – *rs713598*, *rs1726866* and *rs10246939*. Numbers indicate the rounded *D'* and *r*² values, while shading indicates the exact *D'* and *r*² values in regard to the distribution of haplotypes in SNPs

Table 4. Haplotype frequency distribution

Haplotype	Obese group	Control group
CCC	98.96 (49.5)	68.00 (43.6)
GCC	4.00 (2.0)	11.00 (7.1)
GTT	90.96 (45.5)	77.00 (49.4)
CCT	2.04 (1.0)	0.00 (0.0)
GTC	4.04 (2.0)	0.00 (0.0)

Data presented as absolute (relative) percentage (*n* (%)). *N* = 200 for the obese group and *N* = 156 for the control group. Frequency <0.03 was ignored in the analysis.

Obesity, non-taster and the DMFT/dmft score

Among the studied SNPs, the *rs713598* (C-G), *rs1726866* (C-T) and *rs10246939* (C-T) base changes were taken into consideration.^{13,21} Children with the haplotype containing a combination of proline, alanine and valine (PAV) constituted the group highly sensitive to the bitter taste. In contrast, children with the haplotype containing the combination of alanine, valine and isoleucine (AVI) constituted the group with the inability to sense bitterness. Children partially sensitive to the bitter taste (i.e., those with haplotypes PAV/AAV, PAV/AVV) were included in the taster haplotype group (PAV) (Table 5). The mean DMFT/dmft scores for the obesity, non-taster, obese_non-taster, and control_non-taster variables are presented in Table 6. The mean DMFT/dmft score for the study population was 3.017, and of these children, 56.2% were obese and 73.5% were non-tasters. Among non-tasters, obese children constituted 42.7% and children with normal weight constituted 29.8% of the population.

Formula 1, which was created to determine the DMFT/dmft index, depending on the obesity and non-taster dummy variants of the child in the Poisson model analysis, is presented below²¹:

$$DMFT = e^{1.109^{***} - 0.172 * OBESITY + 0.121 NONTASTER} \quad (1)$$

where:

e – constant;

*, ***, – statistical significance level of 10% and 1%, respectively.

Table 5. Distribution of the taster and non-taster groups in control and obese children according to the DMFT/dmft index

Haplotype	Group	n	DMFT/dmft score		p-value
			M ±SD	min–max	
PAV	control taster	53	3.60 ±4.63	0–17	0.120
	obese taster	76	1.96 ±2.09	0–13	
AVI	control non-taster	25	3.15 ±3.59	0–14	0.117
	obese non-taster	24	3.07 ±3.36	0–15	

PAV – proline, alanine and valine; AVI – alanine, valine and isoleucine.

Table 6. Mean DMFT/dmft scores for the variants used in the Poisson and negative binomial models

Variable	DMFT/dmft score	
	M ±SD	min–max
DMFT/dmft	3.017 ±3.507	0–20
Obesity	0.562 ±0.498	0–20
Non-taster	0.735 ±0.448	0–20
Obese_non-taster	0.427 ±0.496	0–20
Control_non-taster	0.298 ±0.459	0–20

A minus sign in front of the coefficient indicates a decrease in DMFT/dmft, while a plus sign indicates an increase.

Obesity, obese non-taster, control non-taster and the DMFT/dmft score

The covariates adjusted according to obesity and the absence of the bitter taste data while constructing the Poisson and negative binomial models for the determination of the DMFT/dmft index are presented in Table 7. Before interpreting the analysis results, the likelihood ratio (LR) test was applied to the 2 competitor models. It was observed that the truncated negative binomial model was more compatible with the data.

Formula 2, which was created to determine the DMFT/dmft index, depending on the obesity, obese_non-taster and control_non-taster variants of the child in the Poisson model analysis, is presented below²¹:

$$DMFT = e^{1.2809^{***} - 0.6088^{***} OBESITY + 0.4482^{**} OBESNONTASTER - 0.1332 CONTROLNONTASTER} \quad (2)$$

where:

e – constant;

*, **, *** – statistical significance level of 10%, 5% and 1%, respectively.

A minus sign in front of the coefficient indicates a decrease in DMFT/dmft, while a plus sign indicates an increase.

According to Formula 1, the DMFT/dmft index was lower in obese children and higher in non-tasters. According to Formula 2, obesity reduces the DMFT/dmft index, although the index was increased in obese non-tasters and decreased in normal-weight non-tasters.

Table 7. Coefficients of the Poisson and negative binomial models for the DMFT/dmft scores

Model	Variable	Poisson	Negative binomial
1 st model	constant	1.109 ±0.094***	1.1619 ±0.2161***
	obesity	–0.172 ±0.087*	–0.2334 ±0.2025
	non-taster	0.121 ±0.010	0.1792 ±0.2269
	AIC	1079.11	802.7377
	SBC	1088.65	815.4649
	expected mean DMFT/dmft	3.017	3.020
2 nd model	constant	1.2809 ±0.1054***	1.3864 ±0.2610***
	obesity	–0.6088 ±0.1800***	–0.7053 ±0.3869
	obese_non-taster	0.4482 ±0.1600**	0.4989 ±0.3236
	control_non-taster	–0.1332 ±0.1300	–0.1728 ±0.3180
	AIC	1072.93	803.0060
	SBC	1085.66	818.9095
	expected mean DMFT/dmft	3.017	3.220

AIC – Akaike information criterion; SBC – Schwarz Bayesian criterion; *, **, *** statistical significance level of 10%, 5% and 1%, respectively.

Discussion

It is important to determine individual risk factors for dental caries in order to create a specific treatment procedure for a particular patient.²² Considering the close association between obesity and the eating habits, researchers have identified a link between obesity and tooth decay.²³ Studies to date have yielded controversial results regarding the relationship between obesity and tooth decay, and further studies are required to understand the link better. A previous systematic review demonstrated that 48% of the investigated studies had reported no significant relationship between BMI and tooth decay; in contrast, a positive and a negative relation was found in 35% and 19% of the same studies, respectively.²⁴ Hong et al. reported a positive relationship between BMI and tooth decay in children aged 60–72 months.²⁵ However, this relationship was not significant when age, genetic predisposition and the eating habits were excluded from the analysis.²⁵ In the present study, a comparison of the DMFT/dmft index for obese and normal-weight children was made, revealing a decreased index value in obese children, and thus hypothesis I was rejected. There are several factors that can be considered responsible for the inverse relation noted between DMFT/dmft and obesity. Children with multiple untreated decayed teeth may be undernourished, as they suffer from severe pain. Also, as a cause of infection, restlessness and sleep problems, tooth decay may decrease life quality in these children, negatively affecting their growth and development. Furthermore, most of the control group subjects were referred to our clinic with a complaint of pain and the control group had a higher incidence of tooth decay than expected, which may have affected our findings.

Many studies have indicated that diet, overweight, low weight, caries, and obesity are interrelated in children. Two recent systematic reviews support this assumption.^{26,27} Therefore, it is reasonable to think that dental caries and obesity may be related to sweet taste genetics. The sugar intake and taste preferences have been associated with the *TAS1R2* polymorphisms in humans. While a high sugar intake has been linked with caries and obesity, few studies have examined the genetic background.^{28,29} Various polymorphism studies of the *TAS1R2* gene have revealed a significant relationship between caries and genotypes.^{29,30} In the present study, the risk of caries was higher in normal-weight children; it may have been due not to the lower incidence of caries in obese children, but to the fact that the population already had a high risk of caries.

A meta-analysis study suggested that *TAS2R38* *rs713598* SNP might play an essential role in caries.³¹ However, our study did not show significant results in any of the 3 SNPs, which is similar to a study by Shimomura-Kuroki et al.³² Since the *TAS2R38* gene

haplotype analysis includes the genotype analysis, the results regarding haplotypes are more critical.³³ The physiology of taste has not been fully clarified to date, although the possible effect of the biological mechanism of taste on tooth decay and obesity appears to be worth investigating.³⁴ The results of the present study demonstrate that tooth decay is associated with 3 different polymorphisms of the *TAS2R38* gene, and the taster and non-taster haplotypes formed by these polymorphisms in obese and normal-weight children. Furthermore, the eating habits and the preferred foods, as the major factors affecting obesity and closely related to tooth decay, may be influenced by bitter taste genetics. There have been few studies to date investigating the relationship between taste genes, obesity and tooth decay, and the conducted ones have failed to clarify the relationship fully.^{14,16,35}

The potential links between taste genetics and various disorders have been investigated in several studies.^{36–38} Accordingly, there is a growing interest in the effects of the biological mechanisms involved in taste sensation on the development of tooth decay and obesity.^{32,38,39} Studies investigating the relationship between the *TAS2R38* polymorphisms and tooth decay reported a lower incidence of tooth decay among tasters. Wendel et al. reported that *TAS2R38* was the main gene associated with tooth decay.⁴⁰ A study conducted on infants revealed that *TAS2R38* and *TAS1R2* controlled their nutritional preferences for the weaning foods.⁴¹ On the other hand, a study conducted on adults argued that this effect might change with age, maternal dietary practices and community habits.⁴² Individuals with the PAV haplotype have greater sensitivity to the bitter taste and better gustatory perception. Due to their ability to sense different flavors, their tendency to cariogenic foods may be reduced.³⁶ In the present study, the participants were classified into tasters and non-tasters, based on the haplotype analysis of the *TAS2R38* gene. Three haplotypes of this gene generated by 3 different SNPs were examined (the corresponding amino acids were A49P, V262A and I296V). In the investigated SNPs, the *rs713598* (C-G), *rs1726866* (C-T) and *rs10246939* (C-T) base exchanges were taken into account. According to the present study results, tooth decay decreases in the presence of obesity and increases in the absence of the bitter taste. However, in the subgroup analysis of non-tasters, the DMFT/dmft index was raised in obese non-tasters and lowered in normal-weight, control non-tasters, and thus hypothesis II was rejected. The authors suggest an increased risk of caries in the obese non-taster group. Further research on this topic can be done. More studies may be needed to explain this issue more clearly. Accordingly, the authors are planning to conduct different studies involving the same patient population. For example, the level of taste can be evaluated by applying solutions of bitter taste and the results can be compared with the genetic analysis.

This study has some limitations, such as keeping the sample size low, and not investigating other causes of caries and obesity.

The topic of taste has not been given the attention it deserves in dentistry or other branches dealing with the sensation of taste. The sensation of taste is a field that is not yet fully understood and there are many gaps in the acquired knowledge. There are branches in medicine that deal directly with the 4 senses of sight (eye), hearing (ear), touch (skin), and smell (nose). However, no specialty field exists for the sense of taste (tongue). The sense of taste and its genetics require support from studies, including the present one.

Conclusions

The CC control homozygous genotype showed higher caries rates in all SNPs of the *TAS2R38* gene, although not statistically significantly. Although there were no statistically significant differences in the DMFT/dmft index with regard to *TAS1R2 rs9701796* SNP in both control and obese groups, a high DMFT/dmft score was observed in the CC homozygous genotype. This may suggest that the CC genotype of *TAS2R38* and *TAS1R2* is prone to caries. Advice including dietary recommendations, increased oral hygiene and controlled calorie intake can be given to children with the identified taste gene polymorphisms.

Ethics approval and consent to participate

This cross-sectional, observational study was approved by the Ethics Committee of the Health Sciences Institute at Atatürk University, Erzurum, Turkey (No. B.30.2.ATA.OA10000-617), and was supported by the Scientific Research Projects Commission at Atatürk University (project No. 2012/66 and 2012/375). The participants' parents provided informed written consent prior to any investigation.

Data availability


The datasets generated and/or analyzed during the current study are available from the corresponding author on reasonable request.


Consent for publication

Not applicable.


ORCID iDs


Münevver Kiliç  <https://orcid.org/0000-0002-3470-0430>

Taskin Gurbuz  <https://orcid.org/0000-0002-8911-6530>

Cigdem Yuce Kahraman  <https://orcid.org/0000-0003-1957-9596>

Atila Cayir  <https://orcid.org/0000-0001-9776-555X>

Abdülbaki Bilgiç  <https://orcid.org/0000-0002-1003-7072>

Yusuf Kurt  <https://orcid.org/0000-0003-3550-1115>

References

- Wang HY, Petersen PE, Bian JY, Zhang BX. The second national survey of oral health status of children and adults in China. *Int Dent J*. 2002;52(4):283–290. doi:10.1111/j.1875-595x.2002.tb00632.x
- Meyer F, Enax J. Early childhood caries: Epidemiology, aetiology, and prevention. *Int J Dent*. 2018;2018:1415873. doi:10.1155/2018/1415873
- Farsi DJ, Elkhodary HM, Merdad LA, et al. Prevalence of obesity in elementary school children and its association with dental caries. *Saudi Med J*. 2016;37(12):1387–1394. doi:10.15537/smj.2016.12.15904
- Honne T, Pentapati K, Kumar N, Acharya S. Relationship between obesity/overweight status, sugar consumption and dental caries among adolescents in South India. *Int J Dent Hyg*. 2012;10(4):240–244. doi:10.1111/j.1601-5037.2011.00534.x
- Satkunam M, Anderson LN, Carsley S, et al. Severe obesity in children 17 to 24 months of age: A cross-sectional study of TARGET Kids! and Better Outcomes Registry & Network (BORN) Ontario. *Can J Public Health*. 2018;109(4):489–497. doi:10.17269/s41997-018-0065-2
- Cinar AB, Christensen LB, Hede B. Clustering of obesity and dental caries with lifestyle factors among Danish adolescents. *Oral Health Prev Dent*. 2011;9(2):123–130. doi:10.3290/j.ohpd.a22012
- Smail HO. The roles of genes in the bitter taste. *AIMS Genet*. 2019;6(4):88–97. doi:10.3934/genet.2019.4.88
- Khan AM, Al-Jandan B, Bugshan A, et al. Correlation of PTC taste status with fungiform papillae count and body mass index in smokers and non-smokers of Eastern Province, Saudi Arabia. *Int J Environ Res Public Health*. 2020;17(16):5792. doi:10.3390/ijerph17165792
- Kim MR, Kusakabe Y, Miura H, Shindo Y, Ninomiya Y, Hino A. Regional expression patterns of taste receptors and gustducin in the mouse tongue. *Biochem Biophys Res Commun*. 2003;312(2):500–506. doi:10.1016/j.bbrc.2003.10.137
- Robertson A, Tirado C, Lobstein T, et al., eds. Food and health in Europe: A new basis for action. World Health Organization. Regional Office for Europe. 2004. <https://apps.who.int/iris/handle/10665/272255>. Accessed June 1, 2021.
- Guo SW, Reed DR. The genetics of phenylthiocarbamide perception. *Ann Hum Biol*. 2001;28(2):111–142. doi:10.1080/03014460151056310
- Jeruzal-Świątecka J, Fendler W, Pietruszewska W. Clinical role of extraoral bitter taste receptors. *Int J Mol Sci*. 2020;21(14):5156. doi:10.3390/ijms21145156
- Drayna D. Human taste genetics. *Annu Rev Genomics Hum Genet*. 2005;6:217–235. doi:10.1146/annurev.genom.6.080604.162340
- Hegde AM, Sharma A. Genetic sensitivity to 6-n-propylthiouracil (PROP) as a screening tool for obesity and dental caries in children. *J Clin Pediatr Dent*. 2008;33(2):107–111. PMID:19358375.
- Lin BP. Caries experience in children with various genetic sensitivity levels to the bitter taste of 6-n-propylthiouracil (PROP): A pilot study. *Pediatr Dent*. 2003;25(1):37–42. PMID:12627700.
- Rupesh S, Nayak UA. Genetic sensitivity to the bitter taste of 6-n-propylthiouracil: A new risk determinant for dental caries in children. *J Indian Soc Pedod Prev Dent*. 2006;24(2):63–68. doi:10.4103/0970-4388.26018
- Kulkarni B, Mattes R. Evidence for presence of nonesterified fatty acids as potential gustatory signaling molecules in humans. *Chem Senses*. 2013;38(2):119–127. doi:10.1093/chemse/bjs095
- Kim UK, Jorgenson E, Coon H, Leppert M, Risch N, Drayna D. Positional cloning of the human quantitative trait locus underlying taste sensitivity to phenylthiocarbamide. *Science*. 2003;299(5610):1221–1225. doi:10.1126/science.1080190
- Bretz WA, Corby PM, Melo MR, et al. Heritability estimates for dental caries and sucrose sweetness preference. *Arch Oral Biol*. 2006;51(12):1156–1160. doi:10.1016/j.archoralbio.2006.06.003
- Neyzi O, Bundak R, Gökçay G, et al. Reference values for weight, height, head circumference, and body mass index in Turkish children. *J Clin Res Pediatr Endocrinol*. 2015;7(4):280–293. doi:10.4274/jcrpe.2183
- Batra M, Shah AF, Rajput P, Shah IA. Comparison of linear and zero-inflated negative binomial regression models for appraisal of risk factors associated with dental caries. *Indian Soc Pedod Prev Dent*. 2016;34(1):71–75. doi:10.4103/0970-4388.175521
- Jepsen S, Blanco J, Buchalla W, et al. Prevention and control of dental caries and periodontal diseases at individual and population level: Consensus report of group 3 of joint EFP/ORCA workshop on the boundaries between caries and periodontal diseases. *J Clin Periodontol*. 2017;44(Suppl 18):S85–S93. doi:10.1111/jcpe.12687

23. Ribeiro Silva AE, Baptista Menezes AM, Demarco FF, Vargas-Ferreira F, Peres MA. Obesity and dental caries: Systematic review. *Rev Saude Publica*. 2013;47(4):799–812. doi:10.1590/S0034-8910.2013047004608
24. Hooley M, Skouteris H, Millar L. The relationship between childhood weight, dental caries and eating practices in children aged 4–8 years in Australia, 2004–2008. *Pediatr Obes*. 2012;7(6):461–470. doi:10.1111/j.2047-6310.2012.00072.x
25. Hong L, Ahmed A, McCunniff M, Overman P, Mathew M. Obesity and dental caries in children aged 2–6 years in the United States: National Health and Nutrition Examination Survey 1999–2002. *J Public Health Dent*. 2008;68(4):227–233. doi:10.1111/j.1752-7325.2008.00083.x
26. Giacaman RA. Sugars and beyond. The role of sugars and the other nutrients and their potential impact on caries. *Oral Dis*. 2018;24(7):1185–1197. doi:10.1111/odi.12778
27. Chi DL, Scott JM. Added sugar and dental caries in children: A scientific update and future steps. *Dent Clin North Am*. 2019;63(1):17–33. doi:10.1016/j.cden.2018.08.003
28. Holla LI, Linhartova PB, Lucanova S, et al. *GLUT2* and *TAS1R2* polymorphisms and susceptibility to dental caries. *Caries Res*. 2015;49(4):417–424. doi:10.1159/000430958
29. Haznedaroğlu E, Koldemir-Gündüz M, Bakır-Coşkun N, et al. Association of sweet taste receptor gene polymorphisms with dental caries experience in school children. *Caries Res*. 2015;49(3):275–281. doi:10.1159/000381426
30. Robino A, Bevilacqua L, Pirastu N, et al. Polymorphisms in sweet taste genes (*TAS1R2* and *GLUT2*), sweet liking, and dental caries prevalence in an adult Italian population. *Genes Nutr*. 2015;10(5):485. doi:10.1007/s12263-015-0485-z
31. Chisini LA, Cademartori MG, Muniz Conde MC, et al. Single nucleotide polymorphisms of taste genes and caries: A systematic review and meta-analysis. *Acta Odontol Scand*. 2021;79(2):147–155. doi:10.1080/00016357.2020.1832253
32. Shimomura-Kuroki J, Nashida T, Miyagawa Y, Sekimoto T. The role of genetic factors in the outbreak mechanism of dental caries. *J Clin Pediatr Dent*. 2018;42(1):32–36. doi:10.17796/1053-4628-42.1.6
33. Melis M, Errigo A, Crnjar R, Pes GM, Barbarossa IT. *TAS2R38* bitter taste receptor and attainment of exceptional longevity. *Sci Rep*. 2019;9(1):18047. doi:10.1038/s41598-019-54604-1
34. Lambert JD, VanDusen SR, Cockroft JE, Smith EC, Greenwood DC, Cade JE. Bitter taste sensitivity, food intake, and risk of malignant cancer in the UK Women's Cohort Study. *Eur J Nutr*. 2019;58(5):2111–2121. doi:10.1007/s00394-018-1772-4
35. Calancie L, Keyserling TC, Taillie LS, et al. *TAS2R38* predisposition to bitter taste associated with differential changes in vegetable intake in response to a community-based dietary intervention. *G3 (Bethesda)*. 2018;8(6):2107–2119. doi:10.1534/g3.118.300547
36. Choi JH, Lee J, Choi IJ, Kim YW, Ryu KW, Kim J. Genetic variation in the *TAS2R38* bitter taste receptor and gastric cancer risk in Koreans. *Sci Rep*. 2016;6:26904. doi:10.1038/srep26904
37. Cossu G, Melis M, Sarchioto M, et al. 6-n-propylthiouracil taste disruption and *TAS2R38* nontasting form in Parkinson's disease. *Mov Disord*. 2018;33(8):1331–1339. doi:10.1002/mds.27391
38. Riccio MP, Franco C, Negri R, et al. Is food refusal in autistic children related to *TAS2R38* genotype? *Autism Res*. 2018;11(3):531–538. doi:10.1002/aur.1912
39. Deshaware S, Singhal R. Genetic variation in bitter taste receptor gene *TAS2R38*, PROP taster status and their association with body mass index and food preferences in Indian population. *Gene*. 2017;627:363–368. doi:10.1016/j.gene.2017.06.047
40. Wendell S, Wang X, Brown M, et al. Taste genes associated with dental caries. *J Dent Res*. 2010;89(11):1198–1202. doi:10.1177/0022034510381502
41. Cont G, Paviotti G, Montico M, et al. *TAS2R38* bitter taste genotype is associated with complementary feeding behavior in infants. *Genes Nutr*. 2019;14:13. doi:10.1186/s12263-019-0640-z
42. Telatar GY, Saydam F, Güzel Aİ, Telatar BC. Variants in taste genes on caries risk and caries activity status. *Med Mol Morphol*. 2020;53(4):244–251. doi:10.1007/s00795-020-00263-5