

Changes in self-reported sleep and awake bruxism in relation to the management of temporomandibular disorders (“care as usual”) in a specialty clinic population

Thiprawee Chattratrai^{1,2,A–F}, Magdalini Thymi^{1,A–F}, Naichuan Su^{3,A,C–F}, Frank Lobbezoo^{1,A,D–F}

¹ Department of Orofacial Pain and Dysfunction, Academic Centre for Dentistry Amsterdam (ACTA), University of Amsterdam and Vrije Universiteit Amsterdam, the Netherlands

² Department of Masticatory Science, Faculty of Dentistry, Mahidol University, Bangkok, Thailand

³ Department of Oral Public Health, Academic Centre for Dentistry Amsterdam (ACTA), University of Amsterdam and Vrije Universiteit Amsterdam, the Netherlands

A – research concept and design; B – collection and/or assembly of data; C – data analysis and interpretation;

D – writing the article; E – critical revision of the article; F – final approval of the article

Dental and Medical Problems, ISSN 1644-387X (print), ISSN 2300-9020 (online)

Dent Med Probl. 2024;61(5):697–704

Address for correspondence

Thiprawee Chattratrai
E-mail: t.chattratrai@acta.nl

Funding sources

Thiprawee Chattratrai was supported with an academic development scholarship by Mahidol University, Bangkok, Thailand.

Conflict of interest

None declared

Acknowledgements

None declared

Received on July 10, 2024

Reviewed on August 27, 2024

Accepted on September 9, 2024

Published online on October 31, 2024

Cite as

Chattratrai T, Thymi M, Su N, Lobbezoo F. Changes in self-reported sleep and awake bruxism in relation to the management of temporomandibular disorders (“care as usual”) in a specialty clinic population. *Dent Med Probl.* 2024;61(5):697–704. doi:10.17219/dmp/193125

DOI

10.17219/dmp/193125

Copyright

Copyright by Author(s)

This is an article distributed under the terms of the

Creative Commons Attribution 3.0 Unported License (CC BY 3.0)

(<https://creativecommons.org/licenses/by/3.0/>).

Abstract

Background. The treatment of temporomandibular disorders (TMD) often includes the management of sleep bruxism (SB) and awake bruxism (AB). However, few studies have investigated how SB and AB change after the initiation of the interventions aimed at reducing the activity of masticatory muscles in TMD patients.

Objectives. The aim of the present study was to investigate changes in self-reported SB and/or AB with regard to baseline at 6 weeks after receiving TMD treatment, i.e., counseling alone or counseling combined with any other treatment, and to investigate the association between the type of TMD treatment and changes in self-reported SB and/or AB.

Material and methods. A total of 68 TMD patients were included in this prospective study, and they all received counseling. Thirty-three of the 68 patients received additional treatment, e.g., physical therapy, psychological therapy and/or an oral appliance, beside counseling. The self-reported SB and AB frequency values were obtained from the Oral Behavior Checklist (OBC) questionnaire at baseline (t_0) and at week 6 after receiving treatment (t_1). The frequency of SB and AB was assessed as SB, AB-grinding, AB-clenching, AB-bracing, and AB-combined (i.e., the maximum frequency of all AB types combined). The Wilcoxon signed-rank test was used to compare the SB and AB frequency at t_0 and t_1 in patients who received counseling alone and those who received counseling combined with other treatment. The χ^2 test was used to investigate the association between the type of TMD treatment and changes in SB and/or AB.

Results. The frequency of self-reported SB and all types of AB did not change in patients who received counseling only. In contrast, there was a significant increase in the frequency of AB-bracing and AB-combined between t_0 and t_1 in patients who received counseling combined with other treatment.

Conclusions. No changes in the frequency of self-reported SB and all types of AB were found in patients who received counseling only. However, patients who received counseling combined with other treatment showed a significant increase in the frequency of AB-bracing and AB-combined as compared to baseline.

Keywords: treatment, follow-up, temporomandibular disorders, sleep bruxism, awake bruxism

Introduction

Sleep bruxism (SB) is a masticatory muscle activity during sleep, characterized by rhythmic or non-rhythmic movement, while awake bruxism (AB) is a repetitive masticatory muscle activity during wakefulness, characterized by tooth contact and/or the bracing or thrusting of the lower jaw.¹ Bruxism is not considered a disorder, but rather a behavior.¹ The prevalence of self-reported SB ranges from 8.0% to 31.4%, while the prevalence of self-reported AB ranges from 22.1% to 31.0% in the general adult population.² Sleep and awake bruxism have been found to be associated with psychosocial factors, such as stress, depression and anxiety.³ Moreover, SB and AB are often investigated for their association with temporomandibular disorders (TMD). The term 'temporomandibular disorders' refers to a group of conditions related to the temporomandibular joint (TMJ), masticatory muscles and associated structures.⁴ The prevalence of TMD symptoms in the adult population is 10.3–30.7%.⁵ Common symptoms of TMD are pain, joint sounds and limited jaw movement.⁴ The TMD pain has been found to be associated with possible and definite AB.⁶ A study found that a higher frequency of self-reported AB, including tooth grinding and clenching, and the bracing of the jaw, was associated with painful TMD.⁷ As for SB, possible SB has been found to be associated with the TMD pain and pain interference with daily life activities,⁸ but the association between definite SB and the TMD pain is inconsistent.^{6,9} A previous study found that probable sleep and awake bruxism, i.e., SB and AB confirmed via a clinical examination, were associated with pain-related TMD.¹⁰ In addition, another study found that 90% of probable sleep bruxers reported jaw-muscle symptoms, such as pain, tiredness or soreness; however, no association was found between muscle activity measured by electromyography (EMG) and jaw-muscle symptoms.¹¹

Temporomandibular disorders constitute a multifactorial condition associated with psychological factors (e.g., stress, depression and anxiety), sleep quality and decreased quality of life (QoL).^{12,13} In addition, the pain and fear related to jaw movements have been associated with the decision to seek care for the TMD pain.¹⁴ The management of TMD includes multidisciplinary non-invasive treatment, such as counseling, physical therapy, medications, and oral appliance therapy. Invasive treatment, such as TMJ surgery, are less common, and only performed in selected cases.^{4,15} The goals of treatment are pain reduction and the recovery of the jaw function.⁴ Given the longstanding notion that SB and AB are viewed as masticatory muscle activities that can overload the masticatory system and contribute to the persistence of the TMD pain, TMD treatment strategies often involve the management of SB and/or AB.^{15–17} Counseling, including education and behavioral modification, can be implemented to reduce AB,¹⁸ and has been shown to reduce the TMD pain and improve the jaw function.^{15,16,18} In addition, the awareness of having AB might help reduce pain.¹⁶ Sleep bruxism is managed through oral appliances, which

aim to reduce the loading of the masticatory system due to the forces exerted while bruxing.¹⁹ Biofeedback treatment has been investigated, as it could reduce a jaw muscle activity during sleep,^{20,21} as well as during wakefulness,¹⁶ but has not yet been implemented as part of routine treatment for the TMD pain.²² Even though SB and AB are common targets in the management of TMD, very few studies have investigated how self-reports of SB and AB change after starting interventions that aim at reducing these masticatory muscle activities in TMD patients.^{18,23}

The present study aimed to investigate changes in self-reported SB and/or AB with regard to baseline at 6 weeks after receiving TMD treatment, i.e., counseling alone or counseling combined with any other treatment, and to investigate the association between the type of TMD treatment and changes in self-reported SB and/or AB. We hypothesized that changes in self-reported SB and/or AB are associated with the type of TMD treatment. More specifically, we hypothesized that counseling combined with any other treatment may alleviate self-reported SB and AB to a greater extent than counseling alone.

Methods

Study sample

A prospective cohort study was performed in the specialty Clinic for Orofacial Pain and Dysfunction of Academic Centre for Dentistry Amsterdam (ACTA), Amsterdam, the Netherlands, from July 2021 until April 2023.

Patients who were referred to the Clinic for Orofacial Pain and Dysfunction of ACTA were eligible to be enrolled in the study if they met the following inclusion criteria:

- at least 18 years old;
- a diagnosis of the TMD pain and/or dysfunction based on the Diagnostic Criteria for Temporomandibular Disorders (DC/TMD),²⁴ for which treatment would be initiated; and
- signed informed consent.

There was no exclusion for medical or dental reasons. Patients who did not complete the online questionnaire at week 6 after starting treatment (t_1) and those who did not receive counseling treatment were excluded.

The study was approved by the ACTA Ethics Committee (ref. No. 2021-64846), and followed the principles of the Declaration of Helsinki.

Study procedures

The study comprised 3 phases: baseline (t_0); treatment; and follow-up (t_1). First, the TMD patients completed a set of questionnaires before their first visit to the clinic. Second, following a clinical examination during the initial visit, the clinicians prescribed treatment based on the DC/TMD diagnosis, relevant comorbidities, patient

preferences, and professional judgment. Last, the patients completed an online questionnaire 6 weeks after the start of treatment. Further details regarding the applied materials and methods are provided below.

Baseline (t_0)

As part of usual care, all patients completed a set of diagnostic questionnaires before their first visit to the clinic. These questionnaires referred to demographic variables, i.e., age and sex, as well as the average facial pain intensity (the Graded Chronic Pain Scale (GCPS) questionnaire),²⁵ depression (the Patient Health Questionnaire-9 (PHQ-9)),²⁶ somatization (the Patient Health Questionnaire-15 (PHQ-15)),²⁷ and anxiety (the Generalized Anxiety Disorder-7 (GAD-7)).²⁸ These questionnaires are part of DC/TMD.²⁴ During the patients' first visit to the clinic, intra- and extraoral inspection, as well as clinical examinations according to DC/TMD were performed. The DC/TMD diagnoses were collected and categorized into 3 categories: pain; dysfunction; and combined pain and dysfunction. The pain category included the DC/TMD diagnoses of local myalgia, myofascial pain, myofascial pain with referral, arthralgia, and headache attributed to TMD. The dysfunction category included the DC/TMD diagnoses of anterior disk displacement with reduction, TMJ subluxation and degenerative joint disease.

Treatment

Each patient received counseling at baseline. The patients received information about their diagnosis and the etiology of their complaints, as well as treatment advice. In addition, the patients could receive one or more other kinds of treatment: physical therapy (including myofeedback, stretching exercises, relaxation, and the self-massage of masticatory muscles); psychological therapy (pain education and a workshop on stress coping); and/or an occlusal splint (a hard occlusal stabilization splint) if the patients reported SB.^{29,30} For the purpose of analysis in this study, the type of treatment was categorized into 2 groups: counseling; and counseling with any other treatment.

Follow-up at 6 weeks after starting treatment (t_1)

Changes in SB and AB after the start of treatment were assessed during the follow-up period by means of a questionnaire containing 11 questions that evaluated 3 domains, namely pain and dysfunction,³¹ patient complaints through the patient-specific approach (PSA),³² together with a complaint improvement question, and the frequency of possible SB and AB.³³ The patients received the questionnaire through e-mail 6 weeks after their initial visit to the clinic.

The frequency of self-reported SB and AB was assessed with the Oral Behavior Checklist (OBC) questions 1, 3,

4, and 6.³³ Self-reported SB was assessed with the OBC question 1, i.e., 'clench or grind teeth when asleep based on any information you may have'. The 5 answer options were: never; <1 night/month; 1–3 nights/month; 1–3 nights/week; and 4–7 nights/week. Self-reported AB was assessed with the OBC items 3, 4 and 6, i.e., 'grind teeth together during waking hours' for the AB-grinding type, 'clench teeth together during waking hours' for the AB-clenching type, and 'hold, tighten or tense muscles without clenching or bringing teeth together' for the AB-bracing type. The answer options ranged from 0 (never) to 4 (always). The highest frequency among these 3 questions was used as the maximum frequency of all self-reported AB types combined, i.e., AB-combined. In this study, changes in self-reported SB and AB between t_0 and t_1 were scored as: 1) not improved, if the self-reported SB or AB frequency at t_1 was higher than or equal to the frequency at t_0 ; or 2) improved, if the self-reported SB or AB frequency at t_1 was lower than the frequency at t_0 .

Sample size calculation

The G*Power 3.1.9.7 software (<https://www.psychologie.hhu.de/arbeitsgruppen/allgemeine-psychologie-und-arbeitspsychologie/gpower>)³⁴ was used to calculate the sample size based on the Wilcoxon signed-rank test. The power of the study was 80%, and the significance level alpha was 0.05. The effect size was set as 0.5, as we assumed a medium size of difference between the 2 groups. A sample size of 35 patients was required.

Statistical analysis

Age, the average facial pain intensity score, and the depression, somatization and anxiety scores were checked for data distribution using the Shapiro–Wilk test. Baseline characteristics, i.e., age, sex, the TMD diagnosis, the average facial pain intensity, and the depression, somatization and anxiety scores, were compared between the 2 treatment groups using the χ^2 test, the Mann–Whitney U test or Fisher's exact test. Differences in the average facial pain intensity and the frequency of self-reported SB and AB, based on the total number of patients, between t_0 and t_1 were compared using the Wilcoxon signed-rank test.

To investigate changes in the frequency of self-reported SB and/or AB between t_0 and t_1 for each type of treatment, the Wilcoxon signed-rank test was used for the patients with counseling alone and separately for those who received counseling combined with other treatment.

To investigate the association between changes in self-reported SB and AB, i.e., improved vs. not improved, on one hand and the type of TMD treatment on the other hand, we used the χ^2 test.

The Castor electronic data capture (EDC) program (Ciwit B.V., Amsterdam, the Netherlands) was used for the collection of study data, and data analysis was

performed with IBM SPSS Statistics for Windows, v. 27.0 (IBM Corp., Armonk, USA). This study complies with the STROBE (Strengthening the Reporting of Observational Studies in Epidemiology) guidelines.

Results

There were 172 patients who met the inclusion criteria at baseline (t_0). Of these, 103 patients who did not complete the online questionnaire at week 6 (t_1) and 1 patient who

Table 1. Distribution of patients according to the provided type of treatment of temporomandibular disorders (TMD)

Type of TMD treatment	N = 68
Counseling only	35 (51.5%)
Counseling combined with other treatment:	33 (48.5%)
– physical therapy	13
– psychological therapy	2
– occlusal splint	8
– splint and physical therapy	3
– splint and psychological therapy	1
– splint and physical therapy and psychological therapy	2
– physical therapy and psychological therapy	1
– physical therapy and GrindCare ^{®a}	1
– physical therapy and BruxApp ^b	1
– medication	1

^a GrindCare[®] – biofeedback device (Medotech, Herlev, Denmark);

^b BruxApp – smartphone application form of ecological momentary assessment (EMA) (WMA Italy, Florence, Italy).

did not receive counseling treatment were excluded. In total, 68 patients were included in this study. There was a significant difference in age between the included and excluded patients ($p = 0.003$). However, there were no significant differences between the included and excluded patients in other baseline characteristics at t_0 : sex ($p = 0.539$); the DC/TMD diagnosis ($p = 0.441$); the average facial pain intensity ($p = 0.406$); depression ($p = 0.979$); somatization ($p = 0.616$); and anxiety ($p = 0.702$). Among the 68 patients, there were no significant differences in the average facial pain intensity and the frequency of self-reported SB and AB between t_0 and t_1 (the average facial pain intensity: $p = 0.076$; SB: $p = 0.781$; AB-combined: $p = 0.180$; AB-grinding: $p = 0.853$; AB-clenching: $p = 0.739$; and AB-bracing: $p = 0.110$). The TMD diagnoses in the TMD-pain group included local myalgia ($n = 5$), myofascial pain ($n = 6$), myofascial pain with referral ($n = 6$), arthralgia ($n = 7$), and headache attributed to TMD ($n = 8$). In the TMD-dysfunction group, the diagnoses included anterior disk displacement with reduction ($n = 6$) and TMJ subluxation ($n = 2$). In the combined group, the diagnoses included local myalgia ($n = 16$), myofascial pain ($n = 9$), myofascial pain with referral ($n = 16$), arthralgia ($n = 26$), headache attributed to TMD ($n = 19$), anterior disk displacement with reduction ($n = 23$), TMJ subluxation ($n = 2$), and degenerative joint disease ($n = 5$). Thirty-five of the 68 patients received counseling only, and 33 patients received counseling and other treatment. The types of TMD treatment are shown in Table 1. There were no differences in baseline characteristics between the patients provided with counseling alone and those who received counseling combined with other treatment (Table 2).

Table 2. Comparison of baseline demographic data between the patients provided with counseling alone ($n = 35$) and those who received counseling combined with other treatment ($n = 33$)

Demographic data	Counseling alone ($n = 35$)	Counseling combined with other treatment ($n = 33$)	Total ($N = 68$)	p -value
Age (18–86 years) $M \pm SD$	48.91 \pm 15.10	46.73 \pm 15.92	47.9 \pm 15.4	0.400 ^b
Sex n (%)				
M	7 (20.0)	4 (12.1)	11 (16.2)	0.378 ^a
F	28 (80.0)	29 (87.9)	57 (83.8)	
DC/TMD diagnosis n (%)				
pain	9 (25.7)	8 (24.2)	17 (25.0)	0.876 ^c
dysfunction	5 (14.3)	3 (9.1)	8 (11.8)	
combined pain and dysfunction	21 (60.0)	22 (66.7)	43 (63.2)	
Average facial pain intensity score at baseline (t_0) Me (IQR)	5 (2–7)	6 (4–7)	5 (3–7)	0.129 ^b
Average facial pain intensity score at week 6 (t_1) Me (IQR)	6 (4–7)	6 (3.5–7)	6 (4–7)	0.700 ^b
Depression score Me (IQR)	5 (3–11)	5 (2–8.5)	5 (2.25–10.75)	0.379 ^b
Somatization score Me (IQR)	9 (6–15)	9 (4–12)	9 (5–14)	0.188 ^b
Anxiety score Me (IQR)	5 (2–8)	3 (1–8)	4 (1–8)	0.206 ^b

M – mean; SD – standard deviation; Me – median; IQR – interquartile range; M – male; F – female; DC/TMD – Diagnostic Criteria for Temporomandibular Disorders²⁴; ^a χ^2 test; ^b Mann–Whitney U test; ^c Fisher's exact test.

Among the patients with counseling alone, the frequency of self-reported SB and all types of self-reported AB did not differ between t_0 and t_1 (Table 3). On the other hand, the frequency of AB-bracing and AB-combined at t_1 was significantly increased among the patients with counseling and other treatment as compared to t_0 (Table 4).

Table 5 shows a significant association between improvement with regard to AB-combined and the type of TMD treatment ($p = 0.023$). Specifically, 78.6% of patients who reported the alleviation of AB-combined and 73.3% of patients who reported the alleviation of AB-bracing were the patients who received counseling alone. In other words, the patients who received counseling

alone were significantly more likely to show improvement than those with the combined treatment.

Discussion

The present study aimed to investigate changes in self-reported SB and/or AB with regard to baseline at 6 weeks after receiving TMD treatment, i.e., counseling alone or counseling combined with any other treatment. The results showed that the frequency of self-reported SB and all types of AB did not change in patients who received counseling only. In contrast, in patients who received counseling combined with other treatment, there was a significant increase in the frequency of AB-bracing and AB-combined between baseline and week 6 after receiving treatment. This may imply that patients who received counseling with other kind of treatment became more aware of the presence of AB-combined and AB-bracing after receiving treatment as compared to baseline.

A previous study found that patients who believed that jaw-overuse behaviors like AB might cause jaw pain tended to report a higher frequency of such behaviors as compared to those who believed that there were other reasons for jaw pain.³⁵ In the present study, 63.6% of patients from the counseling and other treatment group received physical therapy, which could indicate that multiple treatment might increase the awareness of having AB in the patients who received such combined treatment. When patients receive multiple treatment, they may recall and recognize more AB events than they did before treatment. Initially, patients may not be aware of their AB until they receive information about it during counseling. Furthermore, repeated exposure to this information through physical or psychological therapy sessions, for example, may increase patients' awareness of their AB behaviors more than in the case of patients who receive such information only once.

Table 3. Comparisons between the frequency of sleep bruxism (SB) and awake bruxism (AB) at baseline (t_0) and at 6 weeks after receiving treatment (t_1) among the patients with counseling alone ($n = 35$) (Wilcoxon signed-rank test)

Type of bruxism	Frequency at t_0	Frequency at t_1	p -value
SB	4 (0–4)	3 (1–4)	0.188
AB-grinding	0 (0–2)	0 (0–2)	0.748
AB-clenching	3 (1–3)	2 (1–3)	0.527
AB-bracing	2 (1–3)	2 (1–3)	0.472
AB-combined	3 (2–4)	3 (2–3)	0.059

Data presented as median (interquartile range) (Me (IQR)).

Table 4. Comparisons between the frequency of sleep bruxism (SB) and awake bruxism (AB) at baseline (t_0) and at 6 weeks after receiving treatment (t_1) among the patients with counseling and other treatment ($n = 33$) (Wilcoxon signed-rank test)

Type of bruxism	Frequency at t_0	Frequency at t_1	p -value
SB	4 (1.5–4)	3 (1–4)	0.405
AB-grinding	0 (0–2)	0 (0–1)	0.485
AB-clenching	2 (0.5–3)	2 (1–3)	0.255
AB-bracing	2 (0–3)	3 (2–3)	0.008*
AB-combined	2 (1–3)	3 (2–3)	<0.004*

Data presented as Me (IQR).

Table 5. Association between changes in sleep bruxism (SB) and awake bruxism (AB) in terms of improvement and the type of treatment of temporomandibular disorders (TMD) (χ^2 test)

Type of bruxism	Improvement	Counseling alone ($n = 35$)	Counseling combined with other treatment ($n = 33$)	Total ($N = 68$)	p -value
SB	not improved	29 (82.9)	24 (72.7)	53 (77.9)	0.314
	improved	6 (17.1)	9 (27.3)	15 (22.1)	
AB-grinding	not improved	27 (77.1)	26 (78.8)	53 (77.9)	0.870
	improved	8 (22.9)	7 (21.2)	15 (22.1)	
AB-clenching	not improved	22 (62.9)	25 (75.8)	47 (69.1)	0.250
	improved	13 (37.1)	8 (24.2)	21 (30.9)	
AB-bracing	not improved	24 (68.6)	29 (87.9)	53 (77.9)	0.055
	improved	11 (31.4)	4 (12.1)	15 (22.1)	
AB-combined	not improved	24 (68.6)	30 (90.9)	54 (79.4)	0.023*
	improved	11 (31.4)	3 (9.1)	14 (20.6)	

Data presented as number (percentage) (n (%)).

* statistically significant.

Thus, increasing patients' awareness would be beneficial for bruxism management, especially for AB.³⁶ This is in contrast with a previous study finding that counseling and self-management strategies, like self-relaxation, self-massage, stretching exercises, and warm/cold compresses, reduced masticatory muscle pain and AB activity, as measured by surface EMG in female TMD patients after 8 weeks of treatment.³⁷ Meanwhile, usual-care TMD management did not bring improvement with regard to self-reported SB in a brief (6-week) period as compared to self-reported AB. This is in accordance with a previous study showing that sleep hygiene instruction and relaxation techniques did not reduce SB activity, as measured by polysomnography (PSG), when compared between baseline and 4 weeks after the implementation of these techniques.³⁸ It might be difficult for patients to recognize SB events without a report from their sleep partner. However, the present study shows that usual-care TMD treatment can affect self-reported AB in a brief period.

The present study found that there were differences in the frequency of AB-combined and AB-bracing between baseline and week 6 after receiving treatment in patients who received counseling and other treatment. In addition, it was found that after 6 weeks of receiving treatment, 78.6% of patients who reported the alleviation of AB-combined and 73.3% of patients who reported the alleviation of AB-bracing were the patients who received counseling alone. Notwithstanding, there was no significant association between the improvement of AB-bracing and the type of treatment, but, based on the borderline *p*-value, it might have some clinical significance. The percentage of the improvement of AB-combined and AB-bracing in the patients with counseling alone was much higher than in the patients with counseling and other treatment: 31.4% vs. 9.1% for AB-combined; and 31.4% vs. 12.1% for AB-bracing. These different percentages may represent some clinical significance, namely that different types of treatment may be associated with the awareness of having AB and the AB-bracing subtype more than SB and other AB subtypes. In the sample size calculation, we focused on the comparison of patients with counseling alone and patients who received counseling with any other treatment between 2 time points. Thus, we required at least 35 patients in each group. However, we had 33 patients in the counseling and other treatment group, which indicates that the sample size might not be sufficient. This small sample size (i.e., insufficient power) may be one of the reasons why the type of treatment was not statistically significant with regard to the improvement of AB-bracing. On the other hand, the frequency of self-reported SB, AB-grinding and AB-clenching was comparable between the patients who received different types of treatment, and between those who improved or did not improve the abovementioned behaviors. Since patients provided with counseling and any other treatment may increase their awareness of AB, they may improve their

AB behaviors when we continue monitoring over a longer period than 6 weeks. Future research is needed to investigate this matter.

In this study, there was no significant difference in the average facial pain intensity score between baseline and 6 weeks after receiving treatment. In contrast, a study by Donnarumma et al. showed that counseling and self-management strategies could reduce the TMD pain after 8 weeks of receiving treatment, even though the TMD pain was not significantly different between baseline and at 4 weeks of receiving treatment.³⁷ Similarly, 8 weeks of exercise treatment brought the alleviation of the TMD pain.³⁹ Thus, it is suggested that a longer period than 6 weeks is required to observe a reduction in the TMD pain.

Even though the sample size was small, we noticed some changes between the time the patients received their treatment and before the end of treatment. Despite some unexpected trends observed in this study with regard to changes in AB, it is recommended to apply a questionnaire to monitor changes in TMD complaints and oral behaviors in regular care. It is beneficial to have a standardized protocol to monitor SB, AB, TMD, and psychosocial factors along the treatment process, as we are doing in usual care.

The strength of this study is that, first, we assessed self-reported SB and AB at baseline and at 6 weeks after starting treatment. To the best of our knowledge, no study has observed the effect of TMD treatment over a brief period on changes in self-reported SB and AB. A practice-based research network study found that 96% and 46% of dental practitioners considered an occlusal appliance and occlusal adjustment, respectively, as appropriate bruxism management.⁴⁰ The present study may encourage clinicians to incorporate other kinds of treatment, like counseling and physical therapy, for patients. Clinicians should inform patients that they may become more aware of their AB activity after receiving physical therapy, and patients should subsequently alleviate their AB activity. Moreover, we used part of the OBC questionnaire to assess self-reported AB, and not only the maximum frequency of AB, but also different aspects of AB activity, i.e., grinding, clenching and jaw bracing.

Limitations

There are some limitations to this study. First, the frequency of SB and AB was obtained from self-report, whereas the gold standard of SB and AB assessment is PSG for SB and EMG combined with ecological momentary assessment (EMA) for AB.¹ Sleep bruxism is reported more frequently when assessed through self-report than with PSG.⁴¹ Therefore, using EMG or PSG to assess SB and AB is recommended. In the meantime, the Standardized Tool for the Assessment of Bruxism (STAB) has been developed to assess SB and AB. Using STAB is recommended for future research that would focus on

evaluating the bruxism status, the etiology of bruxism and comorbid conditions.⁴² Second, some patients did not fill out all follow-up questionnaires, which were distributed every 6 weeks after the treatment started. Consequently, we had to include only the 1st follow-up questionnaire. Even though a previous study found that biofeedback could reduce SB and AB events, as measured by EMG, in 3 weeks,⁴³ confirming the cause-and-effect relationship between TMD treatment and SB and/or AB self-reported changes may require a longer period of time. Although 6 weeks is a short period for a longitudinal study, it has clinical relevance as a usual duration for follow-up. Third, we did not measure psychosocial factors after receiving treatment, so we could not monitor changes in the psychosocial status, especially in the patients who received psychological treatment, i.e. in 6 out of the 33 participants who were provided with multiple treatment. Last, due to a small sample size, we did not perform a regression analysis to assess the association between the type of TMD treatment and each type of SB and AB. Future research may require a larger sample size, and should include baseline characteristics for adjustment when assessing these associations.

Conclusions

No changes in the frequency of self-reported SB and all types of AB were found in patients who received counseling only. However, patients who received counseling combined with other treatment showed a significant increase in the frequency of AB-bracing and AB-combined as compared to baseline.

Ethics approval and consent to participate

The study was approved by the Ethics Committee at the Academic Centre for Dentistry Amsterdam (ACTA), Amsterdam, the Netherlands (ref. No. 2021-64846). All participants provided signed informed consent.

Data availability

The datasets used and/or analyzed during the current study are available from the corresponding author on reasonable request.

Consent for publication

Not applicable.

ORCID iDs

Thiprawee Chattrattai  <https://orcid.org/0000-0002-3715-1211>
Magdalini Thymi  <https://orcid.org/0000-0002-8544-6495>
Naichuan Su  <https://orcid.org/0000-0001-8034-9410>
Frank Lobbezoo  <https://orcid.org/0000-0001-9877-7640>

References

- Lobbezoo F, Ahlberg J, Raphael KG, et al. International consensus on the assessment of bruxism: Report of a work in progress. *J Oral Rehabil.* 2018;45(11):837–844. doi:10.1111/joor.12663
- Manfredini D, Winocur E, Guarda-Nardini L, Paesani D, Lobbezoo F. Epidemiology of bruxism in adults: A systematic review of the literature. *J Orofac Pain.* 2013;27(2):99–110. doi:10.11607/jop.921
- Fluerăşu MI, Bocşan IC, Ţig IA, Iacob SM, Popa D, Buduru S. The epidemiology of bruxism in relation to psychological factors. *Int J Environ Res Public Health.* 2022;19(2):691. doi:10.3390/ijerph19020691
- de Leeuw R, Klasser GD, eds., American Academy of Orofacial Pain. *Orofacial Pain: Guidelines for Assessment, Diagnosis, and Management.* 6th ed. Batavia, IL: Quintessence Publishing Co., Inc.; 2018:327.
- Iodice G, Cimino R, Vollaro S, Lobbezoo F, Michelotti A. Prevalence of temporomandibular disorder pain, jaw noises and oral behaviours in an adult Italian population sample. *J Oral Rehabil.* 2019;46(8):691–698. doi:10.1111/joor.12803
- Baad-Hansen L, Thymi M, Lobbezoo F, Svensson P. To what extent is bruxism associated with musculoskeletal signs and symptoms? A systematic review. *J Oral Rehabil.* 2019;46(9):845–861. doi:10.1111/joor.12821
- Barbosa C, Manso MC, Reis T, Soares T, Gavinha S, Ohrbach R. Are oral overuse behaviours associated with painful temporomandibular disorders? A cross-sectional study in Portuguese university students. *J Oral Rehabil.* 2021;48(10):1099–1108. doi:10.1111/joor.13226
- Aguilera AB, Lopez LG, Aguilera EB, et al. Relationship between self-reported sleep bruxism and pain in patients with temporomandibular disorders. *J Oral Rehabil.* 2014;41(8):564–572. doi:10.1111/joor.12172
- Manfredini D, Lobbezoo F. Sleep bruxism and temporomandibular disorders: A scoping review of the literature. *J Dent.* 2021;111:103711. doi:10.1016/j.jdent.2021.103711
- Cigdem Karacay B, Sahbaz T. Investigation of the relationship between probable sleep bruxism, awake bruxism and temporomandibular disorders using the Diagnostic Criteria for Temporomandibular Disorders (DC/TMD). *Dent Med Probl.* 2023;60(4):601–608. doi:10.17219/dmp/158926
- Thymi M, Shimada A, Lobbezoo F, Svensson P. Clinical jaw-muscle symptoms in a group of probable sleep bruxers. *J Dent.* 2019;85:81–87. doi:10.1016/j.jdent.2019.05.016
- Pigozzi LB, Pereira DD, Pattussi MP, et al. Quality of life in young and middle age adult temporomandibular disorders patients and asymptomatic subjects: A systematic review and meta-analysis. *Health Qual Life Outcomes.* 2021;19(1):83. doi:10.1186/s12955-021-01727-7
- Yap AU, Cao Y, Zhang MJ, Lei J, Fu KY. Comparison of emotional disturbance, sleep, and life quality in adult patients with painful temporomandibular disorders of different origins. *Clin Oral Investig.* 2021;25(6):4097–4105. doi:10.1007/s00784-020-03740-4
- Rollman A, Visscher CM, Gorter RC, Naeije M. Care seeking for orofacial pain. *J Orofac Pain.* 2012;26(3):206–214. PMID:22838005
- de Barros Pascoal AL, Porto de Freitas RF, Grangeiro da Silva LF, Costa Oliveira AG, Dos Santos Calderon P. Effectiveness of counseling on chronic pain management in patients with temporomandibular disorders. *J Oral Facial Pain Headache.* 2020;34(1):77–82. doi:10.11607/ofph.2163
- Haggiag A, Speciali JG. A new biofeedback approach for the control of awake bruxism and chronic migraine headache: Utilization of an awake posterior interocclusal device. *Arq Neuropsiquiatr.* 2020;78(7):397–402. doi:10.1590/0004-282x20200031
- Noguchi T, Kashiwagi K, Fukuda K. The effectiveness of stabilization appliance therapy among patients with myalgia. *Clin Exp Dent Res.* 2020;6(2):244–253. doi:10.1002/cre2.266
- Xu L, Cai B, Lu S, Fan S, Dai K. The impact of education and physical therapy on oral behaviour in patients with temporomandibular disorder: A preliminary study. *Biomed Res Int.* 2021;2021:6666680. doi:10.1155/2021/6666680
- Gholampour S, Gholampour H, Khanmohammadi H. Finite element analysis of occlusal splint therapy in patients with bruxism. *BMC Oral Health.* 2019;19(1):205. doi:10.1186/s12903-019-0897-z

20. Raphael KG, Janal MN, Sirois DA, Svensson P. Effect of contingent electrical stimulation on masticatory muscle activity and pain in patients with a myofascial temporomandibular disorder and sleep bruxism. *J Orofac Pain*. 2013;27(1):21–31. doi:10.11607/jop.1029
21. Shimada A, Castrillon EE, Svensson P. Revisited relationships between probable sleep bruxism and clinical muscle symptoms. *J Dent*. 2019;82:85–90. doi:10.1016/j.jdent.2019.01.013
22. Lobbezoo F, Aarab G, Ahlers MO, et al. Consensus-based clinical guidelines for ambulatory electromyography and contingent electrical stimulation in sleep bruxism. *J Oral Rehabil*. 2020;47(2):164–169. doi:10.1111/joor.12876
23. Takeuchi-Sato T, Ono Y, Funato M, Sato H, Suganuma T, Baba K. Efficacy of an email-based recording and reminding system for limiting daytime non-functional tooth contact in patients with temporomandibular disorders: A randomized controlled trial. *J Oral Rehabil*. 2020;47(2):158–163. doi:10.1111/joor.12875
24. Schiffman E, Ohrbach R, Truelove E, et al.; International RDC/TMD Consortium Network, International Association for Dental Research, Orofacial Pain Special Interest Group, International Association for the Study of Pain. Diagnostic Criteria for Temporomandibular Disorders (DC/TMD) for Clinical and Research Applications: Recommendations of the International RDC/TMD Consortium Network and Orofacial Pain Special Interest Group. *J Oral Facial Pain Headache*. 2014;28(1):6–27. doi:10.11607/jop.1151
25. Von Korff M, Ormel J, Keefe FJ, Dworkin SF. Grading the severity of chronic pain. *Pain*. 1992;50(2):133–149. doi:10.1016/0304-3959(92)90154-4
26. Kroenke K, Spitzer RL, Williams JB. The PHQ-9: Validity of a brief depression severity measure. *J Gen Intern Med*. 2001;16(9):606–613. doi:10.1046/j.1525-1497.2001.016009606.x
27. Kroenke K, Spitzer RL, Williams JB. The PHQ-15: Validity of a new measure for evaluating the severity of somatic symptoms. *Psychosom Med*. 2002;64(2):258–266. doi:10.1097/00006842-200203000-00008
28. Löwe B, Decker O, Müller S, et al. Validation and standardization of the Generalized Anxiety Disorder Screener (GAD-7) in the general population. *Med Care*. 2008;46(3):266–274. doi:10.1097/MLR.0b013e318160d093
29. Lobbezoo F, van der Zaag J, van Selms MK, Hamburger HL, Naeije M. Principles for the management of bruxism. *J Oral Rehabil*. 2008;35(7):509–523. doi:10.1111/j.1365-2842.2008.01853.x
30. Manfredini D, Colonna A, Bracci A, Lobbezoo F. Bruxism: A summary of current knowledge on aetiology, assessment and management. *Oral Surg*. 2020;13(4):358–370. doi:10.1111/ors.12454
31. Lövgren A, Visscher CM, Häggman-Henrikson B, Lobbezoo F, Marklund S, Wänman A. Validity of three screening questions (3Q/TMD) in relation to the DC/TMD. *J Oral Rehabil*. 2016;43(10):729–736. doi:10.1111/joor.12428
32. Rollman A, Naeije M, Visscher CM. The reproducibility and responsiveness of a patient-specific approach: A new instrument in evaluation of treatment of temporomandibular disorders. *J Orofac Pain*. 2010;24(1):101–105. PMID:20213035.
33. van der Meulen MJ, Lobbezoo F, Aartman IH, Naeije M. Validity of the Oral Behaviours Checklist: Correlations between OBC scores and intensity of facial pain. *J Oral Rehabil*. 2014;41(2):115–121. doi:10.1111/joor.12114
34. Faul F, Erdfelder E, Lang AG, Buchner A. G*Power 3: A flexible statistical power analysis program for the social, behavioral, and biomedical sciences. *Behav Res Methods*. 2007;39(2):175–191. doi:10.3758/bf03193146
35. van Selms MK, Thymi M, Lobbezoo F. Psychological distress and the belief that oral behaviours put a strain on the masticatory system in relation to the self-report of awake bruxism: Four scenarios. *J Oral Rehabil*. 2024;51(1):170–180. doi:10.1111/joor.13460
36. Emodi-Perlman A, Eli I. One year into the COVID-19 pandemic – temporomandibular disorders and bruxism: What we have learned and what we can do to improve our manner of treatment. *Dent Med Probl*. 2021;58(2):215–218. doi:10.17219/dmp/132896
37. Donnarumma V, Michelotti A, Cimino R, Vollaro S, Cioffi I. Short-term effects of a first-line treatment including counseling and self-management strategies on chronic TMD muscle pain and awake bruxism in women. *J Oral Facial Pain Headache*. 2022;36(1):36–48. doi:10.11607/ofph.3037
38. López MV, van Selms MK, van der Zaag J, Hamburger HL, Lobbezoo F. Do sleep hygiene measures and progressive muscle relaxation influence sleep bruxism? Report of a randomised controlled trial. *J Oral Rehabil*. 2015;42(4):259–265. doi:10.1111/joor.12252
39. Uçar İ, Karartı C, Dadalı Y, Özüdoğru A, Okçu M. Masseter muscle thickness and elasticity in bruxism after exercise treatment: A comparison trial. *J Manipulative Physiol Ther*. 2022;45(4):282–289. doi:10.1016/j.jmpt.2022.07.004
40. Mungia R, Lobbezoo F, Funkhouser E, et al.; National Practice-Based Research Network Collaborator Group. Dental practitioner approaches to bruxism: Preliminary findings from the National Dental Practice-Based Research Network. *Cranio*. 2023:1–9. doi:10.1080/08869634.2023.2192173
41. Maluly M, Andersen ML, Dal-Fabbro C, et al. Polysomnographic study of the prevalence of sleep bruxism in a population sample. *J Dent Res*. 2013;92(Suppl 7):975–1035. doi:10.1177/0022034513484328
42. Thymi M, Farzan A, Ahlberg J, Manfredini D, Lobbezoo F. Qualitative suggestions for the further development of the Standardized Tool for the Assessment of Bruxism (STAB). *Dent Med Probl*. 2024;61(3):323–333. doi:10.17219/dmp/183692
43. Sato M, Iizuka T, Watanabe A, et al. Electromyogram biofeedback training for daytime clenching and its effect on sleep bruxism. *J Oral Rehabil*. 2015;42(2):83–89. doi:10.1111/joor.12233